

KEK-TH-806
 hep-th/0203250
 MARCH, 2002

Resummation and Higher Order Renormalization in 4D Quantum Gravity

KEN-JI HAMADA¹

*Institute of Particle and Nuclear Studies,
 High Energy Accelerator Research Organization (KEK),
 Tsukuba, Ibaraki 305-0801, Japan*

Abstract

Higher order renormalization in 4D quantum gravity is carried out using dimensional regularization with great care concerning the conformal-mode dependence. In this regularization, resummation can be automatically carried out without making an assumption like that of David, Distler and Kawai. In this paper we consider a model of 4D quantum gravity coupled to QED. Resummation inevitably implies a four-derivative quantum gravity. The renormalizability is directly checked up to $O(e_r^6)$ and $O(t_r^2)$, where e_r and t_r are the running coupling constants of QED and the traceless gravitational mode. There is no other running coupling constant in our model. The conformal mode is treated exactly, which means it is unrenormalized. It is found that Hathrell's results are included in our results. As a by-product, it is found that a higher-order gravitational correction to the beta function of QED is negative. An advantage of our model is that in the very high-energy regime, it closely resembles exactly solvable 2D quantum gravity. Thus, we can study physical states of 4D quantum gravity in this regime in parallel to those of 2D quantum gravity, which can be described with diffeomorphism invariant composite fields.

PACS: 04.60.-m, 11.10.Gh, 11.25.Pm

¹E-mail address : hamada@post.kek.jp

1 Introduction and Summary

Einstein's general relativity has been extremely well tested, and it has been believed that the graviton exists in a classical description and to obtain a quantum description, it should be quantized [1, 2, 3]. However, the existence of the graviton at the quantum level has not yet been conclusively demonstrated. It is well known that Einstein's theory is unrenormalizable and that the action is unbounded below. Also, it is doubtful whether the usual graviton picture can be applied beyond the Planck mass scale, where the ordinary particle picture seems to become invalid gravitationally.

In an attempt to resolve the problem of renormalizability, a four-derivative gravity [4]–[19] has been investigated, because in this case the gravitational coupling constant becomes dimensionless, and we can avoid the unboundedness problem of the gravitational action. Furthermore, in four-derivative gravity, neither infrared (IR) effects nor ultraviolet (UV) effects can be ignored. This is in accordance with the concept of background-metric independence, because it implies a loss of physical distance. For these reasons, four-derivative gravity is preferred as a quantum theory of gravity that is independent of the background metric [14]–[19].

However, many problems remain to be solved in existing four-derivative gravity models. One is renormalizability. Although these models are renormalizable in the sense of power-counting [5]–[10], this does not imply renormalizability in the true sense. In fact, it has been pointed out that inconsistencies will appear at higher orders [12, 13]. The second problem is that the R^2 action introduced to make the action bounded below is not asymptotically free [9]. The third problem is unitarity [8, 9, 11]. There is an idea regarding unitarity proposed by Tomboulis [8] based on the argument of Lee and Wick [20]. In short, this idea is that gravitational quantum corrections move a ghost pole to a pair of complex poles on the physical sheet. In relativistic theory, it is known that a vertex decaying from real-pole states to such complex-pole states gives measure-zero contributions. Thus, there is no vertex decaying into ghosts in 4D quantum gravity. The proof of unitarity seems, however, nonrigorous. To begin with, it is not understood what constitutes a physical state in 4D quantum gravity, as the strong IR behavior seems to make physical states change. Ghosts might be concealed by the physical state condition, reflecting the background-metric independence. In this paper we investigate these issues, especially the problems of renormalizability and the nature of physical states.

Recently, through studies of 2D quantum gravity [21]–[27], it has come to be understood that all of the problems mentioned above arise because the formulation does not correctly preserve diffeomorphism invariance. In this paper, we show that dimensional regularization [2], carried out *with great care concerning the conformal-mode dependence* [27, 28, 29, 30], is advantageous because it manifestly preserves diffeomorphism invariance and that the resummation process explained in the next section can be automatically carried out without mak-

ing an assumption like that made in the procedure of David, Distler and Kawai (DDK) [23].

Near two dimensions, the D -dimensional action is almost unique, and it is given by the scalar curvature, R . However, near four dimensions, there is an indefiniteness in D -dimensional gravitational actions. To determine the actions here, we need extra conditions other than diffeomorphism invariance. Our proposal is that there are D -dimensional actions, F_D and G_D , which become, respectively, *relevant* and *marginal* at the quantum level, and by using these actions we can construct a renormalizable model described by

$$I = \int d^D x \sqrt{g} \left\{ \frac{1}{t^2} F_D + b G_D + \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \sum_{j=1}^{n_F} i \bar{\psi}_j \not{D} \psi_j - m^2 R + \Lambda \right\}$$

[which is (4.1), below]. The explicit forms of F_D and G_D are presented in Section 3. The F_D term is naturally given by the square of the D -dimensional Weyl tensor, while the marginal term, G_D , which is a combination reduced to the Euler density at $D = 4$, is rather non-trivial. To determine the form of G_D , we consider the resemblance of the 2D quantum gravity action, R , near two dimensions and G_D near four dimensions. Furthermore, considering the integrability condition [32] for conformal anomalies [33]–[37] near four dimensions, we are able to determine G_D .

All bare coupling constants and fields, except for the bare constant, b , and the conformal mode, are renormalized in the usual manner by introducing a renormalization factor, while the conformal mode, which is treated exactly, is not renormalized [18] and b , which is composed of purely pole terms, is treated in a slightly different way. This model has two coupling constants: the QED interaction, e , and the self-interaction of the traceless gravitational mode, t . The running coupling constants are only their renormalized versions, e_r and t_r . The bare constant b is not an independent coupling constant. It is expanded in the coupling constants t and e , and the effective action associated with the $b G_D$ term has a *scale-invariant* form [12]. These points are discussed in Sections 4 and 5. The renormalizability is directly checked up to $O(e_r^6)$ and $O(t_r^2)$ in Section 6. We find that Hathrell's results [29] are contained in our results.

The beta function for the renormalized coupling constant, t_r , is given by

$$\beta_t = - \left(\frac{n_F}{40} + \frac{10}{3} \right) \frac{t_r^3}{(4\pi)^2} - \frac{7n_F}{288} \frac{e_r^2 t_r^3}{(4\pi)^4} + O(t_r^5)$$

[which is (6.5)]. Thus, the theory is asymptotically free with respect to t_r . For the renormalized coupling constant, e_r , we obtain the beta function

$$\beta_e = \frac{4n_F}{3} \frac{e_r^3}{(4\pi)^2} + \left(4n_F - \frac{8}{9} \frac{n_F^2}{b_c} \right) \frac{e_r^5}{(4\pi)^4} + O(e_r^3 t_r^2, e_r t_r^4)$$

[which is (6.23)]. Here, the term proportional to $1/b_c$ is a gravitational correction, and $b_c = 11n_F/360 + 40/9$. Note that the $O(e_r^5)$ term becomes negative for

$n_F \geq 24$. The results for β_t and β_e seem to suggest that all terms in the beta functions related to gravity are negative. Hence, provided that the coefficients of $O(e_r^3 t_r^2)$ and $O(e_r t_r^4)$ in β_e are negative, we can make e_r and t_r sufficiently small asymptotically at very high energies, when the running starts from near the Planck mass scale, where both coupling constants (especially t_r) may be large.

An advantage of this model is that we can exactly prove background-metric independence for the conformal mode in the limit that all running coupling constants vanish [15, 17]. Such a limit may be realized in the very high-energy region. This free field model is completely analogous to 2D quantum gravity. As is well known, 2D quantum gravity can be solved exactly [22, 23], and its physical states can be classified completely using operator formalism [26]. In four dimensions, it is not simple to construct an operator formalism in which composite fields can be treated. In dimensional regularization, however, there is a close relation between them, and for this reason, physical states in 4D quantum gravity can be investigated in analogy to those in 2D quantum gravity, as discussed in Section 7. The last section is devoted to conclusions and discussion.

2 Resummation in Quantum Gravity

In this section we summarize the idea of resummation in 4D quantum gravity. In general, the idea can be applied to any even-dimensional quantum gravity system.

We first give a general argument regarding resummation in *exactly* four dimensions. Diffeomorphism invariant quantum gravity is formally defined by functional integration over the metric,

$$Z = \int \frac{[g^{-1}dg]_g[df]_g}{\text{vol}(\text{diff.})} \exp\{-I(f, g)\}, \quad (2.1)$$

where f is a matter field and I is an invariant action. The measure of the metric is defined by the norm

$$\|g^{-1}dg\|_g^2 = \int d^4x \sqrt{g} \text{tr}(g^{-1}dg)^2, \quad (2.2)$$

where $(g^{-1}dg)^\mu{}_\nu = g^{\mu\lambda}dg_{\lambda\nu}$.

To formulate a well-defined quantum theory of gravity, we must introduce a non-dynamical background metric, $\hat{g}$, and use the measure defined on this background metric, which is obtained by replacing $\sqrt{g}$ in (2.2) with $\sqrt{\hat{g}}$. However, this replacement violates diffeomorphism invariance. Measures defined on $\hat{g}$ are no longer invariant under a general coordinate transformation. To recover diffeomorphism invariance, we must add a *local* action, S . Hence, we have

$$Z = \int \frac{[g^{-1}dg]_{\hat{g}}[df]_{\hat{g}}}{\text{vol}(\text{diff.})} \exp(-S - I). \quad (2.3)$$

This is the idea of DDK [23], which was first developed in two dimensions [24, 25, 26] and then generalized to four dimensions [14]–[19]. The local action S is a four-derivative quantum mechanical quantity that can be considered a contribution induced from the measures. When we quantize gravity, we must quantize this local action again as a tree action and develop a perturbation theory. We here call this procedure *resummation in quantum gravity*.

Resummation works well in two dimensions, where the traceless mode is non-dynamical, so that 2D quantum gravity can be described as a free theory. The idea of DDK can be put into practice using a procedure in which, after assuming the form of the local action, its overall coefficient is determined from the condition that the theory becomes background-metric independent. However, in four dimensions, the traceless mode becomes dynamical, and there is no method to treat it, except for perturbation. The DDK method becomes unclear at higher orders of the perturbation. Thus, we need a more rigorous method to perform resummation without making any assumption.

The following important observation is believed to be true: When we use dimensional regularization, the results are independent of how the measure is chosen, because in this regularization $\delta^D(0) = \int d^D p = 0$. Here, a question arises: Where has the local action S gone? In this paper we show that information regarding S is completely included between D and 4 dimensions. Thus, dimensional regularization carried out with great care concerning the conformal mode dependence may be a manifestly diffeomorphism invariant regularization. This idea was first applied to 2D quantum gravity by Kawai, Kitazawa and Ninomiya [27] and by Hathrell [28, 29] to field theories in curved spacetime.

3 Dimensional Regularization in 4D Quantum Gravity

There is a problem when we apply dimensional regularization to 4D quantum gravity. In *exactly* four dimensions, we usually use the combinations of the curvatures

$$F_4 = R_{\mu\nu\lambda\sigma}R^{\mu\nu\lambda\sigma} - 2R_{\mu\nu}R^{\mu\nu} + \frac{1}{3}R^2, \quad (3.1)$$

$$G_4 = R_{\mu\nu\lambda\sigma}R^{\mu\nu\lambda\sigma} - 4R_{\mu\nu}R^{\mu\nu} + R^2, \quad (3.2)$$

and R^2 , where F_4 and G_4 are the square of the Weyl tensor and the Euler density in four dimensions, respectively. In D dimensions, however, we can generally consider arbitrary combinations of R^2 , $R_{\mu\nu}R^{\mu\nu}$ and $R_{\mu\nu\lambda\sigma}R^{\mu\nu\lambda\sigma}$, which reduce to F_4 and G_4 at $D = 4$. Here, we describe such combinations as F_D and G_D , respectively.

The proposal of this paper is that *the gravitational actions F_D and G_D are almost uniquely determined such that the actions F_D , G_D and R^2 become, respec-*

tively, relevant, marginal (or scale invariant) and irrelevant at the quantum level, and a renormalizable quantum gravity can be constructed using only two actions, F_D and G_D , apart from lower derivative gravitational actions.

In this paper, we consider a coupled system of gravity and QED. In the following, we first discuss the QED sector and then determine F_D by noting the resemblance to the gauge field action. The action G_D is determined by noting the analogy to the action of 2D quantum gravity near two dimensions and also using the integrability condition for conformal anomalies near four dimensions.

3.1 QED sector and F_D

We first study the QED sector in D dimensions. Although the gauge field action is conformally invariant in exactly four dimensions, it is not in D dimensions. Thus, the gauge field action becomes

$$\frac{1}{4} \int d^D x \sqrt{g} F_{\mu\nu} F^{\mu\nu} = \frac{1}{4} \int d^D x \sqrt{\hat{g}} e^{(D-4)\phi} \bar{g}^{\mu\lambda} \bar{g}^{\nu\sigma} F_{\mu\nu} F_{\lambda\sigma}. \quad (3.3)$$

Here, $F_{\mu\nu} = \nabla_\mu A_\nu - \nabla_\nu A_\mu = \partial_\mu A_\nu - \partial_\nu A_\mu$, and the metric is now decomposed as [27]

$$g_{\mu\nu} = e^{2\phi} \bar{g}_{\mu\nu} \quad (3.4)$$

and

$$\bar{g}_{\mu\nu} = (\hat{g} e^{th})_{\mu\nu} = \hat{g}_{\mu\lambda} \left(\delta^\lambda_\nu + t h^\lambda_\nu + \frac{t^2}{2} (h^2)^\lambda_\nu + \dots \right), \quad (3.5)$$

where $h_{\mu\nu} = \hat{g}_{\mu\lambda} h^\lambda_\nu$ and $tr(h) = 0$. The gravitational coupling constant t is introduced only for the traceless mode.

Because the fermion part is conformally invariant in D dimensions, we can remove the conformal-mode dependence by properly rescaling the fermion fields. Hence, there is no direct coupling between the fermions and the conformal mode.

Next, consider the action F_D , which gives the kinetic term of the traceless mode. This action is analogous to the gauge field action. Thus, the most natural choice of F_D is that expressed in terms of the D -dimensional Weyl tensor, $C^\lambda_{\mu\sigma\nu}$, as

$$\begin{aligned} F_D &= C_{\mu\nu\lambda\sigma} C^{\mu\nu\lambda\sigma} \\ &= R_{\mu\nu\lambda\sigma} R^{\mu\nu\lambda\sigma} - \frac{4}{D-2} R_{\mu\nu} R^{\mu\nu} + \frac{2}{(D-1)(D-2)} R^2. \end{aligned} \quad (3.6)$$

Since this satisfies the equation $F_D = e^{-4\phi} \bar{F}_D$, the action becomes

$$\frac{1}{t^2} \int d^D x \sqrt{g} F_D = \frac{1}{t^2} \int d^D x \sqrt{\hat{g}} e^{(D-4)\phi} \bar{F}_D, \quad (3.7)$$

where the action is divided by t^2 to make the kinetic term of the traceless mode independent of t .

3.2 Determination of G_D

Here, we first point out that G_D is an action that reduces to the Euler density at $D = 4$. Hence, there is a resemblance between G_D near four dimensions and the scalar curvature R near two dimensions. In order to determine G_D , we consider this resemblance.

The scalar curvature in D dimensions, which is the only possibility for the action of 2D quantum gravity, is expanded around $D = 2$ as

$$\int d^D x \sqrt{g} R = \sum_{n=0}^{\infty} \frac{(D-2)^n}{n!} S_n^{(2)} \quad (3.8)$$

and

$$S_n^{(2)}(\phi, \bar{g}) = \int d^D x \sqrt{\hat{g}} \phi^n (\bar{\Delta}_2 \phi + \bar{R}) + O(\phi^n), \quad (3.9)$$

where $\Delta_2 = -\square = -\nabla^\lambda \nabla_\lambda$. Note that the action $S_1^{(2)}$ is the well-known Liouville action, or, the local part of the non-local Polyakov action [21]. The action for $n > 1$ is a generalized form of the Liouville action. Its non-local form is discussed in Section 5.

In analogy to 2D quantum gravity, here we look for a four-derivative diffeomorphism invariant action expressed by the following series around $D = 4$:

$$\int d^D x \sqrt{g} G_D = \sum_{n=0}^{\infty} \frac{(D-4)^n}{n!} S_n^{(4)}, \quad (3.10)$$

with

$$S_n^{(4)}(\phi, \bar{g}) = \int d^D x \sqrt{\hat{g}} \phi^n (2\bar{\Delta}_4 \phi + \bar{E}_4) + O(\phi^n). \quad (3.11)$$

Here, the operator Δ_4 and the quantity E_4 are defined by [12]

$$\Delta_4 = \square^2 + 2R^{\mu\nu} \nabla_\mu \nabla_\nu - \frac{2}{3} R \square + \frac{1}{3} \nabla^\mu R \nabla_\mu \quad (3.12)$$

and

$$E_4 = G_4 - \frac{2}{3} \square R. \quad (3.13)$$

In exactly four dimensions, $\sqrt{g} \Delta_4$ becomes a conformally invariant operator for a scalar field, and E_4 satisfies the equation $\sqrt{g} E_4 = \sqrt{\hat{g}} (4\bar{\Delta}_4 \phi + \bar{E}_4)$, which is a four-dimensional generalization of the equation $\sqrt{g} R = \sqrt{\hat{g}} (2\bar{\Delta}_2 \phi + \bar{R})$ at $D = 2$. Thus, the relation between R at $D = 2$ and E_4 at $D = 4$ is apparent.

The first term of the action, $S_1^{(4)}$, is the local part of the Riegert action [12], which is a four-dimensional version of the Polyakov action. The non-local action for $n > 1$ corresponds to a generalization of the Riegert action, which is also discussed in Section 5.

To determine G_D explicitly, here we apply the argument concerning the integrability condition for conformal anomalies given by Bonora, Cotta-Ramusino

and Reina [32]. Consider the general form of the conformal variation of the effective action in D dimensions,

$$\delta_\omega \Gamma = \int d^D x \sqrt{g} \omega \left\{ \eta_1 R_{\mu\nu\lambda\sigma} R^{\mu\nu\lambda\sigma} + \eta_2 R_{\mu\nu} R^{\mu\nu} + \eta_3 R^2 + \eta_4 \square R + \eta_5 F_{\mu\nu} F^{\mu\nu} \right\}. \quad (3.14)$$

Because proper combinations of terms on the r.h.s. are just possibilities for counterterm actions, or bare actions, we wish to reduce the number of coefficients. To this end, we impose the integrability condition $[\delta_{\omega_1}, \delta_{\omega_2}] \Gamma = 0$ in D dimensions, which gives the following relation among the coefficients (see Appendix A):

$$4\eta_1 + D\eta_2 + 4(D-1)\eta_3 + (D-4)\eta_4 = 0. \quad (3.15)$$

In contrast to the case for $D = 4$, we cannot take η_4 itself to be arbitrary in D dimensions, while η_5 remains arbitrary. We can see that the combination of F_D and G_4 satisfies equation (3.15). As another independent combination, we here introduce a D -dimensional extension of what is called the trivial conformal anomaly,

$$M_D = \square R - \frac{D-4}{4(D-1)} R^2, \quad (3.16)$$

though it is no longer trivial in D dimensions. This combination is obtained from the conformal variation of R^2 as $\delta_\omega \int d^D x \sqrt{g} R^2 = -4(D-1) \int d^D x \sqrt{g} \omega M_D$.

Here, we add a condition implied by the analogy to 2D quantum gravity. Specifically, we look for a combination $E_D = G_4 + \eta M_D$ such that $\int d^D x \sqrt{g} E_D = \int d^D x \sqrt{g} (G_4 - \eta \frac{D-4}{4(D-1)} R^2)$ possesses the property of the series (3.10). This condition determines η uniquely as $\eta = -\frac{4(D-3)^2}{(D-1)(D-2)}$. (The details of the calculation are presented in Appendix B.) Hence, we obtain

$$E_D = G_D - \frac{4(D-3)^2}{(D-1)(D-2)} \square R, \quad (3.17)$$

where

$$G_D = G_4 + \frac{(D-3)^2(D-4)}{(D-1)^2(D-2)} R^2. \quad (3.18)$$

Here, E_D is a D -dimensional generalization of E_4 (3.13), and G_D is just what we seek. As in the cases of F_D , G_4 and $F_{\mu\nu} F^{\mu\nu}$, the conformal variation of the spacetime integral of E_D is proportional to E_D itself multiplied by $D-4$, as

$$\delta_\omega \int d^D x \sqrt{g} E_D = (D-4) \int d^D x \sqrt{g} \omega E_D. \quad (3.19)$$

Thus, on the basis of the argument of the integrability condition and the resemblances to gauge theory and 2D quantum gravity, the most natural set of conformal anomalies near four dimensions may be $\{F_D, E_D, M_D, F_{\mu\nu} F^{\mu\nu}\}$. The combinations F_D and E_D (or G_D), as well as conformally invariant matter actions,² represent

²Mass terms are allowed, because they are harmless to UV divergences.

possibilities for the bare actions of 4D quantum gravity near four dimensions. As discussed in Section 6, the higher-order computations of conformal anomalies in curved spacetime carried out by Hathrell [28, 29, 30] strongly support these combinations.

Expanding the spacetime integral of $\sqrt{g}G_D$ around $D = 4$, we obtain

$$\begin{aligned} & \int d^D x \sqrt{g} G_D \\ &= \int d^D x \sqrt{\hat{g}} \left\{ \bar{G}_4 + (D-4) \left(2\phi \bar{\Delta}_4 \phi + \bar{E}_4 \phi + \frac{1}{18} \bar{R}^2 \right) \right. \\ & \quad \left. + \frac{1}{2} (D-4)^2 \left(2\phi^2 \bar{\Delta}_4 \phi + \bar{E}_4 \phi^2 + O(\phi^2) \right) \right. \\ & \quad \left. + O((D-4)^3) \right\}. \end{aligned} \quad (3.20)$$

Hence, the $O(\phi)$ term in $S_1^{(4)}$ is determined to be $\frac{1}{18} \bar{R}^2$.

We close this section by commenting that the actions derived above are almost unique. We have the freedom to change the $O(\phi^n)$ terms in $S_n^{(2)}$ and $S_n^{(4)}$ as $R \rightarrow R + *(D-2)R$ near two dimensions and $G_D \rightarrow G_D + *(D-4)G_D + \natural(D-4)^3 M_D$ near four dimensions, where $*$ and $\natural$ are arbitrary constants. However, as discussed in Section 7, the $O(\phi^n)$ terms would not affect the physical quantities, such as the beta functions and anomalous dimensions. Hence, in this sense, they are almost unique. An ambiguity proportional to F_D can be absorbed into the F_D term.

4 Renormalization

We conjecture that the following quantum gravity model is renormalizable:

$$I = \int d^D x \sqrt{g} \left\{ \frac{1}{t^2} F_D + b G_D + \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \sum_{j=1}^{n_F} i \bar{\psi}_j \not{D} \psi_j - m^2 R + \Lambda \right\}. \quad (4.1)$$

The Dirac operator on general manifolds is defined by $\not{D} = e^{\mu\alpha} \gamma_\alpha D_\mu$, where e_μ^α is the vierbein field satisfying the relations $e_\mu^\alpha e_{\nu\alpha} = g_{\mu\nu}$ and $e_{\mu\alpha} e^\mu_\beta = \delta_{\alpha\beta}$, and the Dirac gamma-matrices on the flat manifold are normalized as $\{\gamma_\alpha, \gamma_\beta\} = -2\delta_{\alpha\beta}$. The covariant derivative for fermions is given by $D_\mu = \partial_\mu + \frac{1}{2} \omega_{\mu\alpha\beta} \Sigma^{\alpha\beta} + ie A_\mu$, where the connection 1-form and the Lorentz generator are defined by $\omega_{\mu\alpha\beta} = e^\nu_\alpha (\partial_\mu e_{\nu\beta} - \Gamma^\lambda_{\mu\nu} e_{\lambda\beta})$ and $\Sigma^{\alpha\beta} = -\frac{1}{4} [\gamma^\alpha, \gamma^\beta]$.

The renormalization is carried out in the usual way, apart from the G_D term, by replacing the bare quantities in I with the renormalized quantities multiplied by the renormalization factors. To make the model finite, we need the following renormalization factors:

$$A_\mu = Z_3^{1/2} A_\mu^r, \quad \psi_j = Z_2^{1/2} \psi_j^r, \quad h_{\mu\nu} = Z_h^{1/2} h_{\mu\nu}^r. \quad (4.2)$$

The coupling constants of QED and the traceless mode are renormalized as

$$e = Z_e e_r, \quad t = Z_t t_r. \quad (4.3)$$

As discussed below, the Ward-Takahashi identity holds even when QED couples with quantum gravity, so that $Z_e = Z_3^{-1/2}$. The important property of this model is that the renormalization factor for the conformal mode is unity;

$$Z_\phi = 1. \quad (4.4)$$

The running coupling constants of this model are only e_r and t_r . The inverse of the gravitational constant, m^2 , and the cosmological constant, Λ , are also renormalized. The renormalization of the G_D term is discussed in Subsection 4.2.

Of course, diffeomorphism invariance does not prohibit us from considering the extended model consisting of $I' = I + \kappa \int d^D x \sqrt{g} R^2$, where $\kappa > 0$. This model might be renormalized in the perturbation of κ with Eq. (4.4) preserved. At the one-loop level, it becomes asymptotically non-free [14]. On the other hand, the integrability condition seems to suggest that κ has to vanish [12, 13]. In any case, our assertion is that the model can be renormalizable at $\kappa = 0$ provided that matter fields are conformally invariant.

As discussed in the previous section, the action in D dimensions has a non-trivial conformal-mode dependence apart from the fermion sector. As mentioned below, *the renormalization procedure is carried out by regarding, in the Laurent expansion of the bare action, the terms with non-negative powers of $D - 4$ to be propagators and vertices and those with negative powers of $D - 4$ to be counterterms.*

4.1 New vertices in the QED sector and F_D

The renormalization factor is given by the Laurent series of poles whose residues are functions of e_r and t_r . To begin with, Z_3 is taken as

$$Z_3 = 1 + \frac{x_1}{D-4} + \frac{x_2}{(D-4)^2} + \cdots. \quad (4.5)$$

The residues are calculated in Section 6. Using this expression, the gauge field action (3.3) is now given by

$$\begin{aligned} & \frac{1}{4} \int d^D x \sqrt{g} F_{\mu\nu} F^{\mu\nu} \\ &= \frac{1}{4} Z_3 \int d^D x e^{(D-4)\phi} \bar{g}^{\mu\lambda} \bar{g}^{\nu\sigma} F_{\mu\nu}^r F_{\lambda\sigma}^r \\ &= \frac{1}{4} \int d^D x \left\{ \left(1 + \frac{x_1}{D-4} + \frac{x_2}{(D-4)^2} + \cdots \right) F_{\mu\nu}^r F_{\mu\nu}^r \right. \end{aligned} \quad (4.6)$$

$$\begin{aligned}
& + \left(D - 4 + x_1 + \frac{x_2}{D-4} + \cdots \right) \phi F_{\mu\nu}^r F_{\mu\nu}^r \\
& + \frac{1}{2} \left((D-4)^2 + (D-4)x_1 + x_2 + \cdots \right) \phi^2 F_{\mu\nu}^r F_{\mu\nu}^r \\
& + \cdots \Big\},
\end{aligned}$$

where we take $\hat{g}_{\mu\nu} = \delta_{\mu\nu}$, and the same lower spacetime indices on the r.h.s. denotes contraction by $\delta_{\mu\nu}$. In the above expression, we omit the interaction terms between the gauge field and the traceless mode, which can be easily derived from the definition.

Similarly, the action of the traceless mode, $\sqrt{g}F_D$, can be expanded in terms of the renormalized quantities using expression (3.7) and the renormalization factors Z_t and Z_h . Although the expression becomes more complicated, the expansion is straightforward. We do not give it here.

Fermion fields are conformally invariant in D dimensions. As mentioned above, we can remove the conformal-mode dependence from the fermion action by replacing the fermion field, ψ , with the rescaled one, $\psi' = e^{\frac{(D-1)}{2}\phi}\psi$, so that $\int d^D x \sqrt{g} i \bar{\psi} \not{D} \psi = \int d^D x \sqrt{\hat{g}} i \bar{\psi}' \not{D} \psi'$. Because in dimensional regularization the result is independent of whether ψ or ψ' is chosen, we quantize ψ' . In the following, we write ψ' as ψ .

The bare fermion action is expanded in the coupling t in the flat background as

$$\begin{aligned}
& \int d^D x i \bar{\psi} \not{D} \psi \\
& = \int d^D x \left\{ i \bar{\psi} \gamma_\mu \partial_\mu \psi - i \frac{t}{4} (\bar{\psi} \gamma_\mu \partial_\nu \psi - \partial_\nu \bar{\psi} \gamma_\mu \psi) h_{\mu\nu} \right. \\
& \quad + i \frac{t^2}{16} (\bar{\psi} \gamma_\mu \partial_\nu \psi - \partial_\nu \bar{\psi} \gamma_\mu \psi) h_{\mu\lambda} h_{\nu\lambda} + i \frac{t^2}{16} \bar{\psi} \gamma_{\mu\nu\lambda} \psi h_{\mu\sigma} \partial_\lambda h_{\nu\sigma} \\
& \quad \left. - e \bar{\psi} \gamma_\mu \psi A_\mu + \frac{et}{2} \bar{\psi} \gamma_\mu \psi A_\nu h_{\mu\nu} - \frac{et^2}{8} \bar{\psi} \gamma_\mu \psi A_\nu h_{\mu\lambda} h_{\nu\lambda} \right\} + O(t^3),
\end{aligned} \tag{4.7}$$

where $\gamma_{\mu\nu\lambda} = \frac{1}{3!}(\gamma_\mu \gamma_\nu \gamma_\lambda + \text{anti-sym.})$. Replacing the bare quantities with the renormalized quantities, we obtain vertices and counterterms, as usual.

4.2 Kinetic term of the conformal mode and new vertices in bG_D

The UV divergences related to the G_D term are renormalized by writing b as

$$b = \sum_{n=1}^{\infty} \frac{b_n}{(D-4)^n}. \tag{4.8}$$

Here, we take $b_0 = 0$, because the tree part, $b_0 \sqrt{g} G_D$, does not have a kinetic term for the gravitational fields, so that it is not a coupling constant. Then, using

expression (3.20), we obtain

$$\begin{aligned}
& b \int d^D x \sqrt{g} G_D \\
&= \int d^D x \left\{ \left(\frac{b_1}{D-4} + \frac{b_2}{(D-4)^2} + \dots \right) \bar{G}_4 \right. \\
&\quad + \left(b_1 + \frac{b_2}{D-4} + \dots \right) \left(2\phi \bar{\Delta}_4 \phi + \bar{E}_4 \phi + \frac{1}{18} \bar{R}^2 \right) \\
&\quad + \frac{1}{2} \left((D-4)b_1 + b_2 + \dots \right) \left(2\phi^2 \bar{\Delta}_4 \phi + \bar{E}_4 \phi^2 + \dots \right) \\
&\quad \left. + \dots \right\}. \tag{4.9}
\end{aligned}$$

We must give some remarks concerning this counterterm. The first is that the kinetic term of the conformal mode is induced by quantum effects, whose overall coefficient is given by the residue of the simple pole, b_1 . The residue of the double pole, b_2 , produces a local counterterm for the kinetic term of the conformal mode. However, this does not imply that Z_ϕ differs from unity. The second is that, even if we take $b_0 \neq 0$, which corresponds to the shift $G_D \rightarrow G_D + *(D-4)G_D$ discussed at the end of Subsection 3.2, the physical quantities are not affected by its value, as discussed in Section 7. The third is that b is a bare quantity, though we expand it in the renormalized coupling constants.

5 Higher-Order Effective Action

In this section, we give brief comments on the renormalized higher-order effective action. Above, we showed that the Laurent expansions of the bare actions using the renormalized quantities produce new local vertices. The overall coefficients of these vertices are related to the UV divergences of higher-order diagrams. The higher-order diagrams also produce finite non-local terms. Thus, the renormalized effective action of quantum gravity, Γ , is given by the sum of the local renormalized vertices in I , described by I_r , and the non-local actions produced by the loop diagrams, W , so that $\Gamma = I_r + W$ [19].

For the gauge sector, the new vertex $\phi^n F_{\mu\nu}^r F_{\mu\nu}^r$ is related to the non-local term $A_r p^2 (\log p^2 / \mu^2)^n A_r$ in momentum space, where the spacetime indices are omitted. The sum of these terms gives the diffeomorphism invariant effective action. Similarly, the new vertex $\phi^n \bar{F}_D^r$ is related to the non-local term $h_r p^4 (\log p^2 / \mu^2)^n h_r$ in momentum space.

Next, consider the effective action related to the counterterm of the Euler density. In two dimensions, the higher-order effective action related to the new vertices, $S_n^{(2)}$, is given by

$$\int d^2 x \sqrt{g} R_r \overbrace{\frac{1}{\Delta_2^r} R_r \frac{1}{\Delta_2^r} R_r \cdots \frac{1}{\Delta_2^r} R_r}^n$$

$$= 2^n 2 \int d^2x \sqrt{\hat{g}} \phi^n (\bar{\Delta}_2^r \phi + \bar{R}_r) + \text{non-local terms.} \quad (5.1)$$

The $n = 1$ action is the well-known non-local Polyakov action [21]. The important property of this action is that it is scale-invariant. The action for $n > 1$ is a natural scale-invariant generalization of the Polyakov action. Here, note that the first term on the r.h.s. is just the first term in the new vertices, $S_n^{(2)}$. The non-local terms can be obtained from higher-order diagrams. The $O(\phi^n)$ term in $S_n^{(2)}$ is necessary to make the theory diffeomorphism invariant, but it would not affect the form of the effective action directly.

Now, we discuss the case of 4D quantum gravity. A four-dimensional version of the Polyakov action was constructed by Riegert [12]. A natural scale-invariant generalization of the Riegert action is

$$\begin{aligned} & \int d^4x \sqrt{g} E_4^r \overbrace{\frac{1}{\Delta_4^r} E_4^r \frac{1}{\Delta_4^r} E_4^r \cdots \frac{1}{\Delta_4^r} E_4^r}^n \\ &= 4^n 2 \int d^4x \sqrt{\hat{g}} \phi^n (2\bar{\Delta}_4^r \phi + \bar{E}_4^r) + \text{non-local terms.} \end{aligned} \quad (5.2)$$

The first term on the r.h.s. is just the first term in the new vertices, $S_n^{(4)}$. This is one of the reasons why we consider G_D to satisfy Eq. (3.10) with (3.11). The non-local terms may be derived from higher-order loop diagrams. This is also investigated in Subsection 6.1 using a concrete expression.

6 Direct Check of Renormalizability

In this section, we directly check the renormalizability of the model proposed in the previous section up to $O(e_r^6)$ and $O(t_r^2)$.

The procedure of gauge fixing for the traceless mode is presented in Appendix C. Here, we employ the Feynman-type gauge. Then, the propagator of the traceless mode, which is described by a spiral line, becomes

$$\frac{D-2}{2(D-3)} \frac{1}{p^4} (I_H)_{\mu\nu, \lambda\sigma}, \quad (6.1)$$

where I_H is the projection operator to the traceless mode (C.9).

The propagator of the conformal mode, which is described by a solid line, is derived from the action $b_1 S_1^{(4)}$ as

$$\frac{(4\pi)^2}{4b_c} \frac{1}{p^4}. \quad (6.2)$$

For later use, we here introduce the coupling-independent constant b_c by

$$b_1 = \frac{1}{(4\pi)^2} \{b_c + O(e_r^2, t_r^2)\}. \quad (6.3)$$

The value of b_c is determined by computing one-loop diagrams.

In this paper we only consider UV divergences, and do not treat IR divergences. Essentially, there is no problem with IR divergences, because of the presence of the Einstein-Hilbert action and the cosmological constant, which play the role of diffeomorphism invariant IR regulators. Technically, we evaluate diagrams by replacing the fourth-order propagator $1/p^4$ with $1/(p^2 + z^2)^2$, where z is a small mass scale.

In the following, we set $D = 4 - 2\epsilon$. For later use, we introduce the dimensionless quantities described by symbols with tildes, such as $\tilde{t} = t\mu^{-\epsilon}$, $\tilde{e} = e\mu^{-\epsilon}$, $\tilde{b} = b\mu^{2\epsilon}$, and so on, where μ is an arbitrary mass scale. Here, we set the dimension of the conformal mode to zero, while the traceless mode is of dimension $\mu^{-\epsilon}$.

6.1 Two- and three-point functions of the traceless mode

We first review the known results for the two- and three-point functions of the traceless mode, which are related to the counterterms of $\sqrt{\hat{g}}\bar{F}_D$ and $\sqrt{\hat{g}}\bar{G}_D$.




Fig. 1. Two- and three-point functions of the traceless mode.

The renormalization factor Z_t is given by

$$Z_t = 1 - \left(\frac{n_F}{80} + \frac{5}{3} \right) \frac{\tilde{t}_r^2}{(4\pi)^2} \frac{1}{\epsilon} - \frac{7n_F}{288} \frac{\tilde{e}_r^2 \tilde{t}_r^2}{(4\pi)^4} \frac{1}{\epsilon} + O(\tilde{t}_r^4). \quad (6.4)$$

The $O(\tilde{t}_r^2)$ term is the sum of one-loop contributions from the QED sector and the gravitational sector. The contributions from the one-loop diagrams with the internal photon, fermion and conformal mode can be calculated from the two-point functions of the traceless mode [Fig. 1-(a)], because Z_h for these diagrams is gauge invariant, so that the relation $Z_t = Z_h^{-1/2}$ is satisfied at this order. They give the contributions $-n_F/80 - 1/40$ [35] for the QED sector and $1/60$ [15] for the conformal mode. However, the contribution to Z_h from the internal traceless mode is, in general, gauge dependent, so that $Z_t \neq Z_h^{-1/2}$. To calculate the contributions from the traceless mode, the background field method [39] is useful. Introducing the background traceless mode as $\hat{g}_{\mu\nu} = (e^{t\hat{h}})_{\mu\nu}$ and calculating the two-point functions with the external h fields replaced by $\hat{h}$, we obtain

$-199/120$ [9] for Z_t , due to the relation $Z_t = Z_{\hat{h}}^{-1/2}$, where $Z_{\hat{h}}$ is the renormalization factor of the background field $\hat{h}_{\mu\nu}$. The $O(e_r^2 t_r^2)$ term is the sum of two-loop contributions from diagrams in which only photons and fermions propagate in the internal lines [38]. Hence, the beta function for the traceless mode coupling constant t_r is

$$\beta_t = -\left(\frac{n_F}{40} + \frac{10}{3}\right) \frac{t_r^3}{(4\pi)^2} - \frac{7n_F}{72} \frac{e_r^2 t_r^3}{(4\pi)^4} + O(t_r^5). \quad (6.5)$$

Thus, this model is asymptotically free with respect to t_r [9, 8].

The residue of the counterterm of $\sqrt{\tilde{g}}\tilde{G}_D$ can be calculated from the three-point functions of gravitational fields. For the simple pole term, it is known to be

$$\tilde{b}_1 = \frac{1}{(4\pi)^2} \left\{ \frac{11n_F}{360} + \frac{40}{9} - \frac{n_F^2}{6} \frac{\tilde{e}_r^4}{(4\pi)^4} + O(\tilde{t}_r^2) \right\}. \quad (6.6)$$

The coupling-independent term in $\tilde{b}_1$ is the sum of the contributions from the QED sector, $(11n_F + 62)/360$ [35], the conformal mode, $-7/90$ [15], and the traceless mode, $87/20$ [9]. From this, the coupling-independent term b_c defined by equation (6.3) is

$$\tilde{b}_c = \frac{11n_F}{360} + \frac{40}{9}. \quad (6.7)$$

The double pole term is known to be [29]

$$\tilde{b}_2 = \frac{1}{(4\pi)^2} \left\{ \frac{2n_F^3}{9} \frac{\tilde{e}_r^6}{(4\pi)^6} + O(\tilde{t}_r^4) \right\}. \quad (6.8)$$

We comment here on the diffeomorphism-invariant effective action. As discussed in Section 5, it is given by the sum of the local action I_r and the finite contribution from loop diagrams W , which contain both local and non-local terms. The one-loop contribution related to the counterterm of $\sqrt{\tilde{g}}\tilde{G}_D$ has the form

$$W_G(\bar{g}_r) = \frac{b_c}{(4\pi)^2} \int d^4x \sqrt{\tilde{g}} \left\{ \frac{1}{8} \bar{E}_4^r \frac{1}{\Delta_4^r} \bar{E}_4^r - \frac{1}{18} \bar{R}_r^2 \right\}. \quad (6.9)$$

This scale-invariant action can be obtained from corrections to the three-point function of the traceless mode. The $\bar{R}_r^2$ term is needed to guarantee that W_G does not have any contributions from two-point diagrams of the traceless mode in the flat background. We can show that in the sum of the action $b_1 S_1^{(4)}$ in I_r and W_G , the non-invariant $\bar{R}_r^2$ terms do cancel out, and the renormalized effective action becomes manifestly diffeomorphism invariant [19]:

$$\frac{b_c}{(4\pi)^2} S_1^{(4)}(\phi, \bar{g}_r) + W_G(\bar{g}_r) = \frac{b_c}{8(4\pi)^2} \int d^4x \sqrt{\tilde{g}} E_4^r \frac{1}{\Delta_4^r} E_4^r. \quad (6.10)$$

This is the Riegert action, or the case of $n = 1$ in (5.2). Thus, only the first term in $S_1^{(4)}$ determines the form of the effective action, as mentioned in Section 5.

6.2 Two-point functions of the conformal mode

Because the properties of the traceless mode are analogous to those of gauge fields, the renormalization procedure for this mode can be carried out straightforwardly. However, diagrams related to the conformal mode are unusual. We study these diagrams in this and following subsections.

The double-pole term of b appears at $O(e_r^6)$ and, at most, $O(t_r^4)$. This implies that there is no counterterm for two-point functions of the conformal mode up to $O(e_r^4)$ and $O(t_r^2)$. This can be seen by direct computation. We first demonstrate the finiteness of the $O(t_r^2)$ correction, which is given by the sum of diagrams (a) and (b) in Fig. 2. The vertices in these diagrams are obtained from the action $b_1 S_1^{(4)}(\phi, \bar{g})$, and they are summarized in Appendix D. We can see that the UV divergences exactly cancel out [18].




Fig. 2. $O(t_r^2)$ diagrams contributing to $\phi - \phi$.

The $O(e_r^2)$ correction is trivially finite. The $O(e_r^4)$ correction is given by the sum of diagrams (a) to (g) in Fig. 3. Here, the wavy line and the dashed line with arrows denote the photon and fermion, respectively. The subdiagram in (e), appearing as a circle with “2” inside, denotes the sum of the two-loop diagrams for the ordinary photon self-energy. We do not here depict counterterm diagrams that subtract the divergences of the subdiagrams. In the following, such counterterm diagrams for subdivergences are always omitted, and we only depict simple-pole counterterm diagrams for direct subtraction, whose residues are related to the residues of double poles in b and Z_3 .

From counting the number of loops and ϵ 's at the vertices, we can easily see that diagrams (a) to (d) in Fig. 3 yield simple poles. Contrastingly, although diagrams (e) to (g) in Fig. 3 potentially have simple-pole divergences, these become finite because of the renormalization properties of QED that the sum of the two-loop diagrams for the photon self-energy has at most a simple pole, and the four-point function of the photon is finite. Thus, we merely compute diagrams (a) to (d) in Fig. 3, and we can see that their sum becomes finite [29].




Fig. 3. $O(e_r^4)$ diagrams contributing to $\phi-\phi$.

Furthermore, using Hathrell's results [29], we can demonstrate the finiteness of the $O(e_r^6)$ two-point functions of the conformal mode in which only photons and fermions propagate in the internal lines (Fig. 4). For the $O(e_r^6)$ diagrams [Fig. 4-(a)] not depicted explicitly, we can see that the sum of the double poles vanishes. The sum of the simple poles does cancel with the direct counterterm of $O(e_r^6)$ [Fig. 4-(b)], which comes from the double-pole term of b , because, as shown by Hathrell, the divergences always appear with the special relation $b_2 = 2c_1$ at $O(e_r^6)$ if we use the set of counterterms bG_4 and cH^2 , where $H = \frac{R}{D-1}$ and c_1 is the residue of the simple pole of c . This relation is naturally realized in our model. From (3.18), the bare action bG_D implies $c_1 = \frac{(D-3)^2}{D-2}b_2 = \frac{1}{2}b_2 + O(D-4)$.



Fig. 4. $O(e_r^6)$ diagrams contributing to $\phi-\phi$.

The most important observation by Hathrell is that this relation is universal, independent of matter fields. Actually, it has been shown that this relation is satisfied in conformally interacting scalar field theory, by Hathrell [28], and also in non-Abelian gauge theory, by Freeman [30]. Our proposal is, in other words, that this special relation also holds for quantized gravity. The fact that Z_ϕ is unity at

$O(t_r^2)$ computed using the vertices in $b_1 S_1^{(4)}$ is a consequence of the combination G_D . Thus, we see that G_D is a universal combination in 4D quantum gravity.

Now, there is no inconsistency between Hathrell's results and the renormalizability of 4D quantum gravity. Although his results have often been cited as evidence for the unrenormalizability of 4D quantum gravity, because it has been believed that R^2 divergences must vanish in order for an effective action to exist [12, 13], this is not true when we use dimensional regularization.

6.3 On $Z_1 = Z_2$

In this subsection, we show that new gravitational interactions do not violate the Ward-Takahashi identity, $Z_1 = Z_2$.

The $O(t_r^2)$ corrections to Z_2 by gravitational fields are given by two diagrams, (a) and (b) in Fig. 5, in which the traceless mode propagates in the internal line. Since fermion fields do not couple with the conformal mode directly, there is no diagram in which the conformal mode propagates in the internal line at the one-loop level. Hence, the sum of the two diagrams in Fig. 5 yields

$$Z_2 - 1 = -\frac{21}{64} \frac{\tilde{t}_r^2}{(4\pi)^2} \frac{1}{\epsilon} \quad (6.11)$$

for all species of fermions in the Feynman-type gauge.





Fig. 5. $O(t_r^2)$ corrections to Z_2 .

In contrast to the case for Z_2 , the $O(t_r^2)$ corrections to Z_1 are given by the seven diagrams (a) to (g) in Fig. 6. Nevertheless, we can show that the Ward-Takahashi identity holds even when QED couples with quantum gravity.






Fig. 6. $O(t_r^2)$ corrections to Z_1 .

The Ward-Takahashi identity is described by an equation in which the QED vertex function, Λ_μ , is equivalent to a derivative of the fermion self-energy, Σ , with respect to the external fermion momentum. Now, the vertex of type $h^n-\bar{\psi}-\psi$ contains derivatives of fermion fields, so that we must take care when determining the momentum dependence at the vertex. Let us first consider diagram (a) in Fig. 5. The fermion self-energy function from this diagram in the momentum space has, apart from an overall coefficient, the form

$$\Sigma^{(a)} = \gamma_\mu \frac{1}{\not{k} + \not{p}} \gamma_\lambda \frac{1}{k^4} (I_H)_{\mu\nu, \lambda\sigma} (k + 2p)_\nu (k + 2p)_\sigma, \quad (6.12)$$

where p is the external fermion momentum and k is the internal traceless-mode momentum. The momentum dependence, $k + 2p$, comes from the vertex of type $h-\bar{\psi}-\psi$. Thus, the derivative of Σ with respect to p acts not only on the fermion propagator but also on the momentum factor at the vertices, and hence

$$\begin{aligned} \frac{\partial}{\partial p_\rho} \Sigma^{(a)} &= -\gamma_\mu \frac{1}{\not{k} + \not{p}} \gamma_\rho \frac{1}{\not{k} + \not{p}} \gamma_\lambda \frac{1}{k^4} (I_H)_{\mu\nu, \lambda\sigma} (k + 2p)_\nu (k + 2p)_\sigma \\ &\quad + \gamma_\mu \frac{1}{\not{k} + \not{p}} \gamma_\lambda \frac{1}{k^4} (I_H)_{\mu\nu, \lambda\sigma} \{2\delta_{\nu\rho} (k + 2p)_\sigma + 2\delta_{\sigma\rho} (k + 2p)_\nu\}. \end{aligned} \quad (6.13)$$

The first term on the r.h.s. of this relation corresponds to the vertex function $\Lambda_\rho^{(a)}$ from diagram (a) in Fig. 6 when evaluated at zero momentum for the external photon. The second is the sum of $\Lambda_\rho^{(b)}$ and $\Lambda_\rho^{(c)}$ from diagrams (b) and (c) in Fig. 6, respectively. Thus, the sum of the vertex functions from (a) to (c) in Fig. 6 is equivalent to the derivative of $\Sigma^{(a)}$ with respect to p . Hence, we can show that Z_2 from diagram (a) in Fig. 5 is equivalent to the sum of Z_1 from diagrams (a) to (c) in Fig. 6.

We can easily show that Z_2 from diagram (b) in Fig. 5 is equivalent to Z_1 from diagram (d) in Fig. 6. In general, the Ward-Takahashi identity holds for all QED vertex functions obtained by attaching an external photon in all possible ways to a fermion line carrying the external charge flow, as in diagrams (a) to (d) in Fig. 6. This is because the vertex $\bar{\psi}\gamma_\mu\psi A_\nu(h^n)_{\mu\nu}$ is given through the minimal substitution $p_\nu \rightarrow p_\nu - eA_\nu$ from the vertex $\bar{\psi}\gamma_\mu p_\nu\psi(h^n)_{\mu\nu}$, apart from vertices of

the type $h^n - \bar{\psi} - \psi$ with a derivative of the traceless mode, which is not affected by the differentiation with respect to the external fermion momentum.

For the diagrams (e) to (g) in Fig. 6, there is no associated fermion self-energy diagram. However, such diagrams do not have divergences, because a vertex of the type $h - A - A$ contains derivatives of the photon fields such that these diagrams vanish when evaluated at zero momentum for the external photon. Thus, we can show $Z_1 = Z_2$ at $O(t_r^2)$. In general, QED vertex corrections with an external photon that stems from the gravitational vertices of types $h^n - A - A$ and $\phi^n - A - A$ vanish when evaluated at zero momentum.

To see that $Z_1 = Z_2$ holds at higher orders, we need to show that all QED vertex corrections obtained by attaching an external photon to internal fermion loops vanish when evaluated at zero momentum for the external photon. To see this, we need to show that Furry's theorem holds in the case including gravitational fields, that is, that a fermion loop with an odd number of QED vertices vanishes even though arbitrary number of the traceless mode is attached on the fermion loop. This can be easily shown by noting that the photon field changes sign under charge conjugation, but the gravitational field does not change sign. Thus, QED vertex corrections including a fermion loop with an odd number of QED vertices, such as diagram (a) in Fig. 7, vanish.

There are diagrams that are not forbidden by the generalized form of Furry's theorem, such as diagram (b) in Fig. 7. However, such diagrams vanish due to a gauge invariance when evaluated at zero momentum for the external photon. Thus, the identity $Z_1 = Z_2$ holds also at higher orders.



Fig. 7. Examples of QED vertex corrections obtained by attaching an external photon to an internal fermion loop.

6.4 Gravitational corrections to Z_3

The gravitational corrections to Z_3 at $O(t_r^2)$ are given by two diagrams, (a) and (b) in Fig. 8. The UV divergences of these two diagrams exactly cancel, so that there is no contribution to Z_3 at $O(t_r^2)$.




Fig. 8. $O(t_r^2)$ corrections to Z_3 .

At higher order, there is a contribution to Z_3 from gravity. Here, we calculate diagrams including an internal conformal mode. Such contributions appear at $O(e_r^4)$. The diagrams are depicted in Fig. 9. They all yield simple poles. Here, diagrams including the ordinary two-loop photon self-energy and the four-photon interaction as a subdiagram are not depicted in Fig. 9, because each sum of such diagrams is finite, as discussed in Subsection 6.2. From a simple counting of the number of loops and ϵ 's at the vertices, diagrams including $\phi^2 - A - A$ vertices and diagrams including two internal conformal-mode lines are trivially finite at $O(e_r^4)$, and therefore they are also omitted.




Fig. 9. $O(e_r^4)$ diagrams yielding simple poles contributing to Z_3 .

We also calculate the residue of the double-pole term of Z_3 at $O(e_r^6)$. It produces a simple pole counterterm of the type $\phi - A - A$, as discussed in Section 4, which is used to show the finiteness of the vertex corrections below. There are only three diagrams contributing to double poles, which are depicted in Fig. 10.




Fig. 10. $O(e_r^6)$ diagrams yielding double poles contributing to Z_3 .

In summary, we obtain

$$Z_3 = 1 - \frac{4n_F}{3} \frac{\tilde{e}_r^2}{(4\pi)^2} \frac{1}{\epsilon} + \left(-2n_F + \frac{8}{27} \frac{n_F^2}{\tilde{b}_c} \right) \frac{\tilde{e}_r^4}{(4\pi)^4} \frac{1}{\epsilon} \\ + \left(-\frac{8n_F^2}{9} + \frac{8}{81} \frac{n_F^3}{\tilde{b}_c} \right) \frac{\tilde{e}_r^6}{(4\pi)^6} \frac{1}{\epsilon^2} + O(\tilde{e}_r^2 \tilde{t}_r^2, \tilde{t}_r^4). \quad (6.14)$$

Here, the simple-pole and the double-pole terms proportional to $1/\tilde{b}_c$ are the contributions from diagrams in Figs. 9 and 10, respectively. The other terms are well-known QED corrections [40].

We now calculate the beta function for the QED coupling constant,

$$\beta_e = \mu \frac{d}{d\mu} \tilde{e}_r. \quad (6.15)$$

The renormalization group equation (RGE) is defined as an equation stipulating that the bare quantities are independent of the arbitrary mass scale, $\mu \frac{d}{d\mu}(\text{bare actions}) = 0$. Thus, we obtain the equation $\mu \frac{d}{d\mu} e = 0$, so that the beta function can be expressed as

$$\beta_e = -\epsilon \tilde{e}_r + \frac{\tilde{e}_r}{2} \frac{\mu}{Z_3} \frac{dZ_3}{d\mu}. \quad (6.16)$$

The bare constant b is treated in a slightly different way. Since $\mu \frac{d}{d\mu} b = 0$, the dimensionless quantity $\tilde{b}$ satisfies $\mu \frac{d}{d\mu} \tilde{b} = 2\epsilon \tilde{b}$. We can re-expand $\tilde{b}$ using the dimensionless bare coupling constants $\tilde{e}$ and $\tilde{t}$ as $\tilde{b} = \sum_n \frac{\tilde{b}'_n}{(D-4)^n}$, where

$$\tilde{b}'_n = \sum_{r,s \geq 0} \tilde{b}'_{c,n}{}^{(r,s)} \tilde{e}^{2r} \tilde{t}^{2s}, \quad (6.17)$$

and hence $\tilde{b}'_{c,1}{}^{(0,0)} = \frac{\tilde{b}_c}{(4\pi)^2}$. The RGE implies that the coefficients $\tilde{b}'_{c,n}{}^{(r,s)}$ are not running coupling constants and that they satisfy the linear equation

$$\mu \frac{d}{d\mu} \tilde{b}'_{c,n}{}^{(r,s)} = 2(1+r+s) \epsilon \tilde{b}'_{c,n}{}^{(r,s)}. \quad (6.18)$$

Thus, if we wish to derive the beta function from information concerning the poles of the renormalization factor, we must consider $\tilde{b}'_{c,n}{}^{(r,s)}$ as unknown constants and use equation (6.18). After deriving finite expressions, we must substitute the values of $\tilde{b}'_{c,n}{}^{(r,s)}$ into them.

We can now calculate the beta function for the QED coupling constant. Here, the related coefficient is only $\tilde{b}_c$, which satisfies $\mu \frac{d}{d\mu} \tilde{b}_c = 2\epsilon \tilde{b}_c$. For the moment, consider the general expression of Z_3 expanded in $\tilde{e}_r$ and the inverse of $\tilde{b}_c$ as

$$Z_3 = 1 + \frac{A_1}{\epsilon} + \frac{A_2}{\epsilon^2} + \cdots + \frac{1}{\tilde{b}_c} \left(\frac{B_1}{\epsilon} + \frac{B_2}{\epsilon^2} + \cdots \right) + \frac{1}{\tilde{b}_c^2} \left(\frac{C_1}{\epsilon} + \frac{C_2}{\epsilon^2} + \cdots \right) + \cdots, \quad (6.19)$$

where

$$\begin{aligned}
A_1 &= \sum_{n \geq 1} A_{1,n} \tilde{e}_r^{2n}, & A_2 &= \sum_{n \geq 3} A_{2,n} \tilde{e}_r^{2n}, \\
B_1 &= \sum_{n \geq 2} B_{1,n} \tilde{e}_r^{2n}, & B_2 &= \sum_{n \geq 3} B_{2,n} \tilde{e}_r^{2n}, \\
C_1 &= \sum_{n \geq 3} C_{1,n} \tilde{e}_r^{2n}, & C_2 &= \sum_{n \geq 4} C_{2,n} \tilde{e}_r^{2n}.
\end{aligned} \tag{6.20}$$

Solving the simultaneous equations (6.15) and (6.16) order by order in ϵ , we obtain the relations between the residues of double poles and simple poles

$$A_{2,3} = -\frac{1}{3}A_{1,1}A_{1,2}, \quad B_{2,3} = -\frac{1}{4}A_{1,1}B_{1,2}, \tag{6.21}$$

and a finite expression for the beta function in the $\epsilon \rightarrow 0$ limit,

$$\beta_e = -A_{1,1}e_r^3 - 2A_{1,2}e_r^5 - 3A_{1,3}e_r^7 - 3B_{1,2}\frac{e_r^5}{b_c} - 4B_{1,3}\frac{e_r^7}{b_c} - 5C_{1,3}\frac{e_r^7}{b_c^2} + \dots \tag{6.22}$$

The residues summarized by (6.14) do satisfy relation (6.21). Note that the coefficient, 3, in front of the $B_{1,2}$ term implies that the diagrams in Fig. 9 are essentially of the three-loop kind. This is reasonable, because the vertex of type $\phi - A - A$ with e_r^2 essentially corresponds to a one-loop diagram. Thus, the treatment for $\tilde{b}$ mentioned above may be considered a trick to count the number of loops correctly.

Thus, we obtain

$$\beta_e = \frac{4n_F}{3} \frac{e_r^3}{(4\pi)^2} + 4n_F \frac{e_r^5}{(4\pi)^4} - \frac{8}{9} \frac{n_F^2}{b_c} \frac{e_r^5}{(4\pi)^4} + O(e_r^3 t_r^2, e_r t_r^4). \tag{6.23}$$

Substituting the value of b_c into (6.7), we can see that the sign of the $O(e_r^5)$ term becomes negative for $n_F \geq 24$. This result, as well as the result for β_t , (6.5), suggests that all gravitational corrections give negative contributions to the beta functions. Hence, provided that the residues of the simple poles of Z_3 at $O(e_r^2 t_r^2)$ and $O(t_r^4)$ are positive, which give negative contributions to β_e , we can consider the scenario in which the UV fixed point of the QED coupling constant becomes zero, or close to zero, asymptotically beyond the Planck mass scale.

6.5 $O(e_r^6)$ renormalization of $\phi - A - A$ vertex I

The renormalization factor Z_3 has double poles at $O(e_r^6)$. This means that a new counterterm for the vertex of type $\phi - A - A$ appears at $O(e_r^6)$. In the following two subsections, we directly check whether this vertex is renormalizable using only the data for Z_3 and $Z_\phi = 1$.

In this subsection, we first consider the case in which there are no internal gravitational fields. The diagrams that potentially have UV divergences at $O(e_r^6)$ are divided into four groups, as depicted in Fig. 11. These are described in the following.




Fig. 11. $O(e_r^6)$ $\phi-A-A$ vertex diagrams without internal gravitational fields.

Diagram (a) in Fig. 11 represents the sum of diagrams given by attaching a line with e_r^2 at the end to internal photons in the ordinary two-loop photon self-energy (2LPSE) diagrams [Fig. 11-(a)]. Since the sum of the diagrams for 2LPSE gives only a simple pole, diagram Fig.11-(a) also gives a simple pole.




Fig. 11-(a). $\phi-A-A$ vertex diagrams (a).

Diagram (b) in Fig. 11 represents the sum of diagrams given by attaching a line with ϵ at the end to internal photons in the ordinary three-loop photon self-energy (3LPSE) with a single fermion loop [Fig. 11-(b)]. It is known that the sum of the diagrams for 3LPSE with a single fermion loop yields only a simple pole, and thus diagram Fig. 11-(b) becomes finite, due to the ϵ at the vertex.




Fig. 11-(b). $\phi-A-A$ vertex diagrams (b).

Diagram (c) in Fig. 11 represents the sum of diagrams given by attaching a line with ϵ at the end to internal photons in 3LPSE with two fermion loops [Fig. 11-(c)]. Since the sum of the diagrams for 3LPSE with two fermion loops gives a double pole, but not a triple pole, diagram Fig. 11-(c) yields a simple pole.




Fig. 11-(c). $\phi-A-A$ vertex diagrams (c).

Diagram (d) in Fig. 11 is the $O(e_r^6)$ direct counterterm mentioned above. The sum of all UV divergences evaluated at zero momentum for the external conformal mode vanishes exactly:

$$\Gamma_{\mu\nu}^{\phi AA}(0; k, -k)|_I = \left\{ -\frac{8}{3} + \frac{16}{9} + \frac{8}{9} \right\} n_F^2 \frac{e_r^6}{(4\pi)^6} \frac{1}{\epsilon} (\delta_{\mu\nu} k^2 - k_\mu k_\nu) = 0. \quad (6.24)$$

We here normalize the effective action in the momentum representation as $\Gamma = \int [dp][dk][dl] (2\pi)^D \delta^D(p+k+l) \phi(p) A_\mu^r(k) A_\nu^r(l) \Gamma_{\mu\nu}^{\phi AA}(p; k, l)$, where $[dp] = \frac{d^D p}{(2\pi)^D}$. The first and second terms in the braces result from the diagrams in Figs. 11-(a) and 11-(c), respectively. The third term in (6.24) comes from the direct counterterm diagram (d) in Fig. 11 related to the double pole term in Z_3 .

6.6 $O(e_r^6)$ renormalization of $\phi-A-A$ vertex II

Next, we consider the vertex diagrams of type $\phi-A-A$, including internal conformal-mode lines that yield divergences proportional to e_r^6/b_c . The diagrams are depicted in Fig. 12, where, for diagrams (e) to (g), the other assignments of ϵ and e_r^2 on the vertices should be taken into account. As mentioned above, the sum of diagrams at $O(e_r^6)$ including 2LPSE as a subdiagram is finite. The same conclusion holds for diagrams including 3LPSE, four- and six-photon interactions as a subdiagram. Therefore, none of them are depicted in Fig. 12. The new features that do not appear in the previous cases are that there are contributions from vertices of types ϕ^3 and ϕ^2-A-A .




Fig. 12. $O(e_r^6)$ ϕ - A - A vertex diagrams with internal conformal modes.

From a direct calculation, we can see that the sum of the UV divergences from diagrams (d) to (i) in Fig. 12 exactly vanishes when evaluated at zero momentum for the external conformal mode. The sum of the UV divergences from all other diagrams also vanishes:

$$\Gamma_{\mu\nu}^{\phi AA}(0, k, -k)|_{II} = \left\{ -\frac{8}{81} + \frac{16}{81} - \frac{8}{81} \right\} \frac{n_F^3}{b_c} \frac{e_r^6}{(4\pi)^6} \frac{1}{\epsilon} (\delta_{\mu\nu} k^2 - k_\mu k_\nu) = 0. \quad (6.25)$$

The first value in the braces results from the sum of (a) to (c), the second from the sum of (j) to (m), and the third from the direct counterterm diagram (n), whose value is derived from the residue of the double pole of Z_3 .

7 Physical States in 4D Quantum Gravity

That a theory possesses quantum diffeomorphism invariance implies that the theory is independent of the background metric. Thus, in the quantum gravity phase, the physical distance measured by the metric field would become meaningless. This implies that IR effects and UV effects are sufficiently strong. Therefore, the usual Fock states based on the vanishing of correlations among them in the IR limit are no longer physical states in quantum gravity.

The physical state conditions reflecting background-metric independence are represented by the equation $\frac{\delta Z}{\delta \hat{g}^{\mu\nu}} = \langle \hat{T}_{\mu\nu} \rangle = 0$. These are usually called the Hamiltonian momentum constraints. In 4D quantum gravity, we use perturbation theory to treat the traceless mode so that the background-metric independence is slightly violated. This violation would, in principle, vanish if the perturbative calculation were carried out to infinitely high order. However, as in 2D quantum gravity [23, 25], we can prove rigorously the background-metric independence for the conformal mode in the free field limit, $t_r = e_r = 0$ [15, 17], which is realized at very high energies. This is an advantage of our formalism. We can study the structure of the physical states in the very high-energy regime using this free field model.

The physical state conditions are exactly solved in 2D quantum gravity using conformal field theory [22, 23]. Recall that although 2D quantum gravity is a free theory, it has rich structure: an infinite number of physical states [26], a non-linear property of factorization [24], and so on. For instance, the physical states of 2D quantum gravity coupled to a scalar field X with central charge $c_X = 1$ are given by $\int d^2x : e^{2\phi} :$, $\int d^2x : \partial X \bar{\partial} X :$, $\int d^2x : (\partial^2 X \pm 2i(\partial X)^2)(\bar{\partial}^2 X \pm 2i(\bar{\partial} X)^2) e^{-\phi \mp iX} :$, and so on, where $::$ denotes the normal order, and the conformal mode is properly normalized. These infinitely numerous states are called “discrete states.” Here, note that there is an analogy between this two-dimensional field X and the traceless mode h^μ_ν in 4D quantum gravity, because $2n$ -th order fields in $2n$ dimensions have the same IR and UV behavior for every n . As for the conformal mode, the relation to 2D quantum gravity is more apparent: the first term

in the action, $S_1(\phi, \bar{g})$, in each dimension satisfies the Wess-Zumino consistency condition, which is used to prove the background-metric independence for the conformal mode [25, 15, 17]. Thus, although 4D quantum gravity is asymptotically free, the physical states are non-trivial even in the very high-energy regime, which can be described with diffeomorphism invariant composite fields.

One of the simplest physical states is the cosmological constant. We here investigate it for $t_r = e_r = 0$ in detail. In dimensional regularization, the cosmological constant is given by $\Lambda \int d^D x e^{D\phi} = \Lambda \sum_n \int d^D x \frac{D^n}{n!} \phi^n$ in the flat background,³ which is renormalized by replacing the bare constant Λ by $Z_\Lambda \Lambda_r$. Since there is no coupling constant for the conformal mode, we, for the present, evaluate the renormalization factor Z_Λ for large n_F .

The diagrams yielding simple poles at $O(1/b_c)$ and $O(1/b_c^2)$ are given by (a) and (b) in Fig. 13, respectively. Contrastingly, diagrams with separated ovals, for example, diagrams (a) and (b), in Fig. 14 at $O(1/b_c^2)$, do not produce simple poles. More precisely, the sum of the UV divergences from diagram (a), itself, in Fig. 14 and from the associated subdivergence counterterm diagram not depicted here, yields only a double pole, while diagram (b) in Fig. 14 gives no contribution to Z_Λ , because the UV divergence from diagram (b) itself cancels that from its associated subdivergence counterterm diagram. In general, because a diagram with more than two separated ovals does not produce a simple pole, it does not contribute to the anomalous dimension. Thus, we obtain

$$Z_\Lambda = 1 - \frac{2}{\bar{b}_c} \frac{1}{\epsilon} - \frac{2}{\bar{b}_c^2} \frac{1}{\epsilon} + \frac{2}{\bar{b}_c^2} \frac{1}{\epsilon^2} + \cdots, \quad (7.1)$$

and hence the anomalous dimension of the cosmological constant is

$$\gamma_\Lambda = \frac{\mu}{Z_\Lambda} \frac{dZ_\Lambda}{d\mu} = \frac{4}{b_c} + \frac{8}{b_c^2} + \cdots. \quad (7.2)$$




Fig. 13. $O(1/b_c)$ and $O(1/b_c^2)$ corrections to the cosmological constant.

³In previous papers [14, 18], in analogy with the operator formalism of DDK, the anomalous dimension was computed using a procedure in which the deformed cosmological constant field $e^{\alpha\phi}$ is considered, and the equation $\alpha = 4 + \gamma_\Lambda$ is solved for α , where γ_Λ is given by a function of α computed with one-loop diagrams. However, this procedure is actually incorrect in dimensional regularization, because diffeomorphism invariance is obviously violated, though the derived anomalous dimension for the cosmological constant seems correct. Here, we revise the procedure in a manifestly invariant way.





Fig. 14. $O(1/b_c^2)$ corrections to the cosmological constant that do not yield simple poles.

Let us now see what is obtained when summing up all orders. Using the analogy to 2D quantum gravity, the anomalous dimension defined by subtracting the canonical dimension from the quantum dimension was conjectured to be [14, 18]⁴

$$\begin{aligned}\gamma_\Lambda &= 2b_c \left(1 - \sqrt{1 - \frac{4}{b_c}}\right) - 4 \\ &= \frac{4}{b_c} + \frac{8}{b_c^2} + \frac{20}{b_c^3} + \dots\end{aligned}\tag{7.3}$$

The first two terms in the expansion agree exactly with the computed terms. The corresponding diagrams at $O(1/b_c^m)$ are given by diagrams with a single oval in Fig. 15. Here, the first diagram has $m - 1$ ϕ^3 -vertices with ϵb_1 , the second one has $m - 3$ ϕ^3 vertices with ϵb_1 and a ϕ^4 -vertex with $\epsilon^2 b_1$, and so on. The last one is constructed by using a ϕ^{m+1} -vertex with $\epsilon^{m-1} b_1$. Each diagram has m loops multiplied by ϵ^{m-1} , so that it yields a simple pole. Since the other potentially divergent diagrams with multiple ovals do not produce simple poles, only the diagrams in Fig. 15 contribute to the anomalous dimension at $O(1/b_c^m)$.



⁴In two dimensions, the anomalous dimension can be exactly calculated using conformal field theory as $\gamma_\Lambda = \alpha - 2$, where $\alpha = b_c(1 - \sqrt{1 - 4/b_c})$ is the Liouville charge of the cosmological constant operator, : $e^{\alpha\phi}$: [22, 23]. For 2D quantum gravity coupled to N scalar fields, $b_c = (25 - N)/6$. In dimensional regularization, the first two terms of γ_Λ in the expansion w.r.t. $1/b_c$ is also obtained by evaluating the same diagrams in Fig. 13 when using the propagator and the vertices obtained from the bare action, $\frac{b_1}{D-2} \int d^D x \sqrt{g} R$ and the series (3.8), where $b_1 = \frac{b_c}{4\pi}$ can be determined by computing one-loop divergences for the counterterm of $\sqrt{g} \hat{R}$.

Fig. 15. $O(1/b_c^m)$ corrections to the cosmological constant.

Here, there is an important observation concerning the ambiguity of the action G_D . This action is defined by the series (3.10). All vertices used in Fig. 15 stem from the term $\phi^n \bar{\Delta}_4 \phi$ in $S_n^{(4)}$. From a simple counting of the number of loops and ϵ 's at the vertices, we can easily see that, if we use the vertices coming from the $O(\phi^n)$ term in $S_n^{(4)}$, diagrams with a single oval become finite. Also, if we use vertices obtained when $b_0 \neq 0$ in (4.8), the same conclusion is reached. Furthermore, although there is a potentially divergent diagram with the $O(\phi^n)$ vertices in the case that it has more than two separated ovals, it does not yield a simple pole, as mentioned above. Thus, the anomalous dimensions of the cosmological constant can be determined only from the first term in $S_n^{(4)}$.

In this paper, although we have only considered the case of the cosmological constant, generalization is straightforward. It seems natural to assume that the physical state is given by a combination of diffeomorphism invariant composite fields. Thus, we conjecture that the physical state can be classified according to the condition of renormalizability. In that case, all physical quantities, such as anomalous dimensions, would be independent of the choice of the $O(\phi^n)$ term in $S_n^{(4)}$, as in the case of the cosmological constant. In this sense, G_D is almost unique.

8 Discussion

In this paper, we studied the problems of higher-order renormalization and physical states in 4D quantum gravity. Using dimensional regularization with great care concerning the conformal-mode dependence, we carried out the renormalization of a model of 4D quantum gravity coupled to QED up to $O(e_r^6)$ and $O(t_r^2)$. The procedure is manifestly diffeomorphism invariant, and resummation was done without making an DDK-like assumption. As a by-product, we found that higher-order gravitational corrections to the beta functions of e_r and t_r are negative.

This model suggests that a configuration characterized by the vanishing of the Weyl tensor becomes dominant at very high energies, where the coupling t_r is small. Such a configuration is called conformally flat, or $S^4 \times (\text{conformal factor})$. Quantum gravity may be realized in the early universe and on inside of black holes, where spacetime singularities would appear in the classical description. However, this model suggests that such singularities disappear through the effect of quantum gravity [41].

Physical states in our model were discussed in analogy with those in 2D quantum gravity. In particular, the cosmological constant term was studied in detail. In general, even though the theory is asymptotically free, the physical states might be given by diffeomorphism-invariant composite fields, such as glueball states in 4D Yang-Mills theory, because of the strong IR effects. Therefore, to

distinguish them from ordinary gravitons, we here call them “graviball states.” A photon may also be confined as a photonball state in quantum gravity. On the other hand, usual graviton and photon states are low-energy states, which would appear when the background-metric independence, namely the condition $\hat{T}_{\mu\nu}|phys\rangle = 0$, is violated. It is not, at present, understood how to connect the quantum gravity phase with our present Einstein phase. It seems that there is a discontinuity between them.

Finally, we comment on recent interesting numerical results concerning the dynamical triangulation (DT) approach [42, 43, 44] in four dimensions [45]–[49]. It is believed that the DT method realizes the quantum gravity phase in the very high-energy regime. The entropy factor in the DT approach corresponds to the four-derivative gravitational actions, because they are dimensionless in four dimensions, so that they essentially represent quantum effects. The anomalous dimension of the cosmological constant in this regime is almost given by a function of b_c , as discussed in Section 7, and corrections that depend on the coupling constant t_r are small [18]. If we consider a gravity system coupled to n_X conformal scalar fields and n_A gauge fields, b_c is given by a function of $n_X + 62n_A$, and hence γ_Λ is a function of $n_X + 62n_A$. This number dependence should be numerically checked using the DT approach [48].

Acknowledgements

This work was supported in part by a Grant-in-Aid for Scientific Research from the Ministry of Education, Science and Culture of Japan.

Appendix

A Conformal-mode Dependence of Curvatures

The conformal mode dependence of curvatures in D dimensions is obtained by using the expression for the Christoffel symbol $\Gamma_{\mu\nu}^\lambda = \bar{\Gamma}_{\mu\nu}^\lambda + \bar{g}_\mu^\lambda \bar{\nabla}_\nu \phi + \bar{g}_\nu^\lambda \bar{\nabla}_\mu \phi - \bar{g}_{\mu\nu} \bar{\nabla}^\lambda \phi$, as

$$R_{\mu\sigma\nu}^\lambda = \bar{R}_{\mu\sigma\nu}^\lambda + \bar{g}_\nu^\lambda \bar{\Delta}_{\mu\sigma} - \bar{g}_\sigma^\lambda \bar{\Delta}_{\mu\nu} + \bar{g}_{\mu\sigma} \bar{\Delta}_\nu^\lambda - \bar{g}_{\mu\nu} \bar{\Delta}_\sigma^\lambda + (\bar{g}_\nu^\lambda \bar{g}_{\mu\sigma} - \bar{g}_\sigma^\lambda \bar{g}_{\mu\nu}) \bar{\nabla}_\delta \phi \bar{\nabla}^\delta \phi, \quad (\text{A.1})$$

$$R_{\mu\nu} = \bar{R}_{\mu\nu} - (D-2) \bar{\Delta}_{\mu\nu} - \bar{g}_{\mu\nu} \{ \bar{\square} \phi + (D-2) \bar{\nabla}_\lambda \phi \bar{\nabla}^\lambda \phi \}, \quad (\text{A.2})$$

$$R = e^{-2\phi} \{ \bar{R} - 2(D-1) \bar{\square} \phi - (D-1)(D-2) \bar{\nabla}_\lambda \phi \bar{\nabla}^\lambda \phi \}, \quad (\text{A.3})$$

where $\bar{\Delta}_{\mu\nu} = \bar{\nabla}_\mu \bar{\nabla}_\nu \phi - \bar{\nabla}_\mu \phi \bar{\nabla}_\nu \phi$. Our curvature conventions are $R_{\mu\nu} = R_{\mu\lambda\nu}^\lambda$ and $R_{\mu\sigma\nu}^\lambda = \partial_\sigma \Gamma_{\mu\nu}^\lambda - \dots$.

The variation formulae with respect to the conformal change $\delta_\omega g_{\mu\nu} = 2\omega g_{\mu\nu}$ are given by

$$\delta_\omega \sqrt{g} R = (D-2)\omega \sqrt{g} R - 2(D-1)\sqrt{g} \square \omega \quad (\text{A.4})$$

and

$$\delta_\omega \sqrt{g} R^{\mu\nu\lambda\sigma} R_{\mu\nu\lambda\sigma} = (D-4)\omega \sqrt{g} R^{\mu\nu\lambda\sigma} R_{\mu\nu\lambda\sigma} - 8\sqrt{g} R^{\mu\nu} \nabla_\mu \nabla_\nu \omega, \quad (\text{A.5})$$

$$\begin{aligned} \delta_\omega \sqrt{g} R^{\mu\nu} R_{\mu\nu} &= (D-4)\omega \sqrt{g} R^{\mu\nu} R_{\mu\nu} - 2\sqrt{g} R \square \omega \\ &\quad - 2(D-2)\sqrt{g} R^{\mu\nu} \nabla_\mu \nabla_\nu \omega, \end{aligned} \quad (\text{A.6})$$

$$\delta_\omega \sqrt{g} R^2 = (D-4)\omega \sqrt{g} R^2 - 4(D-1)\sqrt{g} R \square \omega, \quad (\text{A.7})$$

$$\begin{aligned} \delta_\omega \sqrt{g} \square R &= (D-4)\omega \sqrt{g} \square R + (D-6)\sqrt{g} \nabla^\lambda R \nabla_\lambda \omega \\ &\quad - 2\sqrt{g} R \square \omega - 2(D-1)\sqrt{g} \square^2 \omega, \end{aligned} \quad (\text{A.8})$$

$$\delta_\omega \sqrt{g} F_{\mu\nu} F^{\mu\nu} = (D-4)\omega \sqrt{g} F_{\mu\nu} F^{\mu\nu}. \quad (\text{A.9})$$

Applying these variation formulae to (3.14), we obtain

$$\begin{aligned} [\delta_{\omega_1}, \delta_{\omega_2}] \Gamma &= \{4\eta_1 + D\eta_2 + 4(D-1)\eta_3 + (D-4)\eta_4\} \\ &\quad \times \int d^D x \sqrt{g} R (\omega_1 \square \omega_2 - \omega_2 \square \omega_1). \end{aligned} \quad (\text{A.10})$$

Thus, the integrability condition is given by (3.15).

B Expansions of Curvatures

From expression (A.3), we can see that the spacetime integral of the scalar curvature is expanded around $D = 2$ as follows:

$$\int d^D x \sqrt{g} R = \sum_{n=0}^{\infty} \frac{(D-2)^2}{n!} \int d^D x \sqrt{\hat{g}} \{ -(D-1)\phi^n \bar{\square} \phi + \bar{R} \phi^n \}. \quad (\text{B.1})$$

This is the action of 2D quantum gravity near two dimensions.

Next, we consider the expansion of four-derivative actions near four dimensions. The spacetime integral of M_D is expanded around $D = 4$ as

$$\begin{aligned} &\int d^D x \sqrt{g} M_D \\ &= -\frac{D-4}{4(D-1)} \int d^D x \sqrt{g} R^2 \\ &= -\frac{1}{4(D-1)} \sum_{n=0}^{\infty} \frac{(D-4)^n}{n!} \int d^D x \sqrt{\hat{g}} \left\{ (D-4)\phi^n \bar{R}^2 - 2(D-1)(D-6)\phi^n \bar{R} \bar{\square} \phi \right. \\ &\quad + 2(D-1)(D-2)\phi^n \bar{\nabla}^\lambda \bar{R} \bar{\nabla}_\lambda \phi + 4(D-1)^2 \phi^n \bar{\square}^2 \phi \\ &\quad + 8(D-1)^2 (D-4)\phi^n \bar{\square} \phi \bar{\nabla}^\lambda \phi \bar{\nabla}_\lambda \phi \\ &\quad \left. + (D-1)^2 (D-2)^2 (D-4)\phi^n (\bar{\nabla}^\lambda \phi \bar{\nabla}_\lambda \phi)^2 \right\}, \end{aligned} \quad (\text{B.2})$$

and, also, the spacetime integral of G_4 is expanded around $D = 4$ as

$$\begin{aligned}
& \int d^D x \sqrt{g} G_4 \\
&= \sum_{n=0}^{\infty} \frac{(D-4)^n}{n!} \int d^D x \sqrt{\hat{g}} \left\{ \phi^n \bar{G}_4 + 4(D-3) \phi^n \bar{R}^{\mu\nu} \bar{\nabla}_\mu \bar{\nabla}_\nu \phi \right. \\
&\quad - 2(D-3) \phi^n \bar{R} \bar{\square} \phi - 2(D-2)(D-3)(D-4) \phi^n \bar{\square} \phi \bar{\nabla}^\lambda \phi \bar{\nabla}_\lambda \phi \\
&\quad \left. - (D-2)(D-3)^2(D-4) \phi^n (\bar{\nabla}^\lambda \phi \bar{\nabla}_\lambda \phi)^2 \right\}. \tag{B.3}
\end{aligned}$$

The last two terms in each of the series in (B.2) and (B.3) give contributions of $O(\phi^{n+2})$ and $O(\phi^{n+3})$ at order $(D-4)^n$, while the series in (3.11), which we seek, gives a contribution of at most $O(\phi^{n+1})$. Thus, in order to obtain the series in (3.11), we must consider the sum of these two actions, $\int d^D x \sqrt{g} (G_4 + \eta M_D)$, and choose the constant η properly in order to remove the last two terms. Although we cannot exactly remove these terms simultaneously, the series we seek can be obtained by merely removing the last term, so that $\eta = -\frac{4(D-3)^2}{(D-1)(D-2)}$. In this way, we obtain

$$\begin{aligned}
& \int d^D x \sqrt{g} G_D \\
&= \sum_{n=0}^{\infty} \frac{(D-4)^n}{n!} \int d^D x \sqrt{\hat{g}} \left\{ \phi^n \bar{E}_D + \frac{4(D-3)^2}{D-2} \phi^n \bar{\square}^2 \phi + 4(D-3) \phi^n \bar{R}^{\mu\nu} \bar{\nabla}_\mu \bar{\nabla}_\nu \phi \right. \\
&\quad - \frac{4(D-3)(D^2-6D+10)}{(D-1)(D-2)} \phi^n \bar{R} \bar{\square} \phi - \frac{2(D-3)^2(D-6)}{(D-1)(D-2)} \phi^n \bar{\nabla}^\lambda \bar{R} \bar{\nabla}_\lambda \phi \\
&\quad \left. - \frac{2(D-3)(D-4)^3}{D-2} \phi^n \bar{\square} \phi \bar{\nabla}_\lambda \phi \bar{\nabla}^\lambda \phi \right\} \\
&= \int d^D x \sqrt{\hat{g}} \left\{ \bar{G}_4 + (D-4) \left(2\phi \bar{\Delta}_4 \phi + \bar{E}_4 \phi + \frac{1}{18} \bar{R}^2 \right) \right. \\
&\quad + \frac{1}{2} (D-4)^2 \left(2\phi^2 \bar{\Delta}_4 \phi + \bar{E}_4 \phi^2 \right. \\
&\quad \quad + 6\phi \bar{\square}^2 \phi + 8\phi \bar{R}^{\mu\nu} \bar{\nabla}_\mu \bar{\nabla}_\nu \phi - \frac{28}{9} \phi \bar{R} \bar{\square} \phi + \frac{8}{9} \phi \bar{\nabla}^\lambda \bar{R} \bar{\nabla}_\lambda \phi \\
&\quad \quad \left. - \frac{14}{9} \bar{R} \bar{\square} \phi + \frac{1}{9} \bar{R}^2 \phi + \frac{5}{54} \bar{R}^2 \right) \\
&\quad + \frac{1}{3!} (D-4)^3 \left(2\phi^3 \bar{\Delta}_4 \phi + \bar{E}_4 \phi^3 - 6 \bar{\square} \phi \bar{\nabla}^\lambda \phi \bar{\nabla}_\lambda \phi + 9\phi^2 \bar{\square}^2 \phi + \dots \right) \\
&\quad \left. + O((D-4)^4) \right\}. \tag{B.4}
\end{aligned}$$

C BRST Formulation

The general coordinate transformations $\delta_\xi g_{\mu\nu} = g_{\mu\lambda} \nabla_\nu \xi^\lambda + g_{\nu\lambda} \nabla_\mu \xi^\lambda$ can be completely decomposed into transformations of the conformal mode and the

traceless mode as

$$\begin{aligned}\delta_\xi \bar{g}_{\mu\nu} &= \bar{g}_{\mu\lambda} \bar{\nabla}_\nu \xi^\lambda + \bar{g}_{\nu\lambda} \bar{\nabla}_\mu \xi^\lambda - \frac{2}{D} \bar{g}_{\mu\nu} \hat{\nabla}_\lambda \xi^\lambda, \\ \delta_\xi \phi &= \xi^\lambda \partial_\lambda \phi + \frac{1}{D} \hat{\nabla}_\lambda \xi^\lambda,\end{aligned}\tag{C.1}$$

where $\bar{\nabla}_\lambda \xi^\lambda = \hat{\nabla}_\lambda \xi^\lambda$ is used. The traceless-mode part can be expanded in the coupling constant t . The general coordinate transformation for A_μ is given by

$$\delta_\xi A_\mu = \xi^\nu \nabla_\nu A_\mu + A_\nu \nabla_\mu \xi^\nu = \xi^\nu \partial_\nu A_\mu + A_\nu \partial_\mu \xi^\nu,\tag{C.2}$$

and the $U(1)$ gauge transformation is $\delta_\lambda A_\mu = \partial_\mu \lambda$.

Let us fix the coordinate transformation invariance for the traceless mode and the gauge invariance of the photon field according to the procedure of the BRST quantization. The BRST transformations can be obtained by replacing ξ^μ/t in the equations for general coordinate transformations, (C.1) and (C.2), and λ in the $U(1)$ gauge transformation with corresponding Grassmann ghost fields, c^μ and c , respectively. We obtain

$$\begin{aligned}\delta_{\mathbf{B}} h^\mu{}_\nu &= \left\{ \hat{\nabla}^\mu c_\nu + \hat{\nabla}_\nu c^\mu - \frac{2}{D} \delta^\mu{}_\nu \hat{\nabla}_\lambda c^\lambda + t c^\lambda \hat{\nabla}_\lambda h^\mu{}_\nu \right. \\ &\quad \left. + \frac{t}{2} h^\mu{}_\lambda (\hat{\nabla}_\nu c^\lambda - \hat{\nabla}^\lambda c_\nu) + \frac{t}{2} h^\lambda{}_\nu (\hat{\nabla}^\mu c_\lambda - \hat{\nabla}_\lambda c^\mu) + \dots \right\}, \\ \delta_{\mathbf{B}} \phi &= t c^\mu \partial_\mu \phi + \frac{t}{D} \hat{\nabla}_\mu c^\mu, \\ \delta_{\mathbf{B}} A_\mu &= \partial_\mu c + t c^\nu \partial_\nu A_\mu + t A_\nu \partial_\mu c^\nu, \\ \delta_{\mathbf{B}} c^\mu &= t c^\nu \hat{\nabla}_\nu c^\mu = t c^\nu \partial_\nu c^\mu, \\ \delta_{\mathbf{B}} c &= t c^\mu \partial_\mu c, \\ \delta_{\mathbf{B}} \tilde{c}^\mu &= B^\mu, \quad \delta_{\mathbf{B}} B^\mu = 0, \\ \delta_{\mathbf{B}} \tilde{c} &= B, \quad \delta_{\mathbf{B}} B = 0,\end{aligned}\tag{C.3}$$

where $c_\mu = \hat{g}_{\mu\nu} c^\nu$. We can easily see that this BRST transformation is nilpotent. Using this transformation, the gauge-fixing term and the FP ghost action can be written as

$$\begin{aligned}I_{GF+FP} &= \int d^D x \sqrt{\hat{g}} \delta_{\mathbf{B}} \left\{ \tilde{c}^\mu \hat{N}_{\mu\nu} \left(\chi^\nu - \frac{\zeta}{2} B^\nu \right) + \tilde{c} \left(\hat{\nabla}^\mu A_\mu - \frac{\alpha}{2} B \right) \right\} \\ &= \int d^D x \sqrt{\hat{g}} \left\{ B^\mu \hat{N}_{\mu\nu} \chi^\nu - \frac{\zeta}{2} B^\mu \hat{N}_{\mu\nu} B^\nu - \tilde{c}^\mu \hat{N}_{\mu\nu} \hat{\nabla}^\lambda \delta_{\mathbf{B}} h^\nu{}_\lambda \right. \\ &\quad \left. + B \hat{\nabla}^\mu A_\mu - \frac{\alpha}{2} B^2 - \tilde{c} \hat{\nabla}^\mu \delta_{\mathbf{B}} A_\mu \right\},\end{aligned}\tag{C.4}$$

where $\chi^\nu = \hat{\nabla}^\lambda h^\nu{}_\lambda$ and $\hat{N}_{\mu\nu}$ is a symmetric second-order operator defined on $\hat{g}_{\mu\nu}$. This action has a covariant form with respect to $\hat{g}_{\mu\nu}$, but not with respect to the

dynamical gravitational modes, ϕ and h^μ_ν . Note that the conformal mode does not couple with all ghost fields, and the ghosts for the $U(1)$ gauge invariance are not coupled to the traceless mode either. If B_μ and B are integrated out, we obtain [9]

$$I_{GF} = \int d^D x \sqrt{\hat{g}} \left\{ \frac{1}{2\zeta} \chi^\mu \hat{N}_{\mu\nu} \chi^\nu + \frac{1}{2\alpha} (\hat{\nabla}^\mu A_\mu)^2 \right\}. \quad (\text{C.5})$$

In the following, we use the flat background, in which $\hat{g}_{\mu\nu} = \delta_{\mu\nu}$. The bilinear term of the traceless mode is given by

$$\frac{1}{t^2} \int d^D x \bar{F}_D = \frac{D-3}{D-2} \left(h_{\mu\nu} \partial^4 h_{\mu\nu} + 2\chi_\mu \partial^2 \chi_\mu - \frac{D-2}{D-1} \chi_\mu \partial_\mu \partial_\nu \chi_\nu \right) + O(t), \quad (\text{C.6})$$

where $\chi_\mu = \partial_\lambda h_{\mu\lambda}$ and $\partial^2 = \partial_\lambda \partial_\lambda$. From this expression, we should choose the following form for the gauge-fixing term:

$$\hat{N}_{\mu\nu} = \frac{2(D-3)}{D-2} \left(-2\partial^2 \delta_{\mu\nu} + \frac{D-2}{D-1} \partial_\mu \partial_\nu \right). \quad (\text{C.7})$$

From expression (C.5), $\zeta = 1$ corresponds to the Feynman-type gauge, in which the gauge-fixing term cancels out the last two of the kinetic terms in (C.6). Then, the propagator of the traceless mode becomes

$$\frac{D-2}{2(D-3)} \frac{1}{p^4} (I_H)_{\mu\nu, \lambda\sigma}, \quad (\text{C.8})$$

where I_H is the projection operator to the traceless mode,

$$(I_H)_{\mu\nu, \lambda\sigma} = \frac{1}{2} \delta_{\mu\lambda} \delta_{\nu\sigma} + \frac{1}{2} \delta_{\mu\sigma} \delta_{\nu\lambda} - \frac{1}{D} \delta_{\mu\nu} \delta_{\lambda\sigma}, \quad (\text{C.9})$$

which satisfies $I_H^2 = I_H$.

D Vertices in $b_1 S_1^{(4)}(\phi, \bar{g})$

From the expansion in (B.4), we find

$$S_1^{(4)}(\phi, \bar{g}) = \int d^D x \sqrt{\hat{g}} \left\{ 2\phi \bar{\Delta}_4 \phi + \bar{E}_4 \phi + \frac{1}{18} \bar{R}^2 \right\}. \quad (\text{D.1})$$

The bare action $b_1 S_1^{(4)}(\phi, \bar{g})$ is expanded in the coupling t in the flat background, apart from the kinetic term of ϕ , as

$$\mathcal{L}_{S_1}^2 = -\frac{2}{3} b_1 t \partial^2 \phi \partial_\mu \partial_\nu h_{\mu\nu} + \frac{b_1}{18} t^2 (\partial_\mu \partial_\nu h_{\mu\nu})^2, \quad (\text{D.2})$$

$$\begin{aligned} \mathcal{L}_{S_1}^3 = 2b_1 t \left(2\partial_\mu \phi \partial_\nu \partial^2 \phi + \frac{4}{3} \partial_\mu \partial_\lambda \phi \partial_\nu \partial_\lambda \phi \right. \\ \left. - \frac{2}{3} \partial_\lambda \phi \partial_\mu \partial_\nu \partial_\lambda \phi - 2\partial_\mu \partial_\nu \phi \partial^2 \phi \right) h_{\mu\nu}, \end{aligned} \quad (\text{D.3})$$

$$\begin{aligned} \mathcal{L}_{S_1}^4 = 2b_1 t^2 \left(\partial^2 \phi \partial_\mu \partial_\nu \phi h_{\mu\lambda} h_{\nu\lambda} + \partial_\mu \partial_\nu \phi \partial_\lambda \partial_\sigma \phi h_{\mu\nu} h_{\lambda\sigma} \right. \\ \left. + \text{terms including } \partial h \right), \end{aligned} \quad (\text{D.4})$$

where $\mathcal{L}_{S_1}^2$ is derived from the term $-\frac{2}{3}b_1(\bar{\square}\bar{R})\phi + \frac{1}{8}b_1\bar{R}^2$. The vertices $\mathcal{L}_{S_1}^3$ and $\mathcal{L}_{S_1}^4$ are from $2b_1\phi\bar{\Delta}_4\phi$.

References

- [1] B. DeWitt, in Relativity, Groups and Topology, eds. B. DeWitt and C. DeWitt (Gordon and Breach, New York, 1964); Phys. Rev. **160** (1967) 1113; Phys. Rev. **162** (1967) 1195, 1239.
- [2] G. 't Hooft and M. Veltmann, Ann. Inst. Henri Poincare **XX** (1974) 69; M. Veltmann, in Methods in Field Theory, Les Houches 1975.
- [3] S. Weinberg, in General Relativity, an Einstein Centenary Survey, eds. S. Hawking and W. Israel (Cambridge Univ. Press, Cambridge, 1979).
- [4] R. Utiyama and B. DeWitt, J. Math. Phys. **3** (1962) 608.
- [5] S. Deser, Proceedings of the Conference on Gauge Theories and Modern Field Theories, edited by R. Arnowitt and P. Nath (MIT Press, Cambridge, 1975); S. Weinberg, Proceedings of the XVIIth International Conference on High Energy Physics, edited by J. R. Smith (Rutherford Laboratory, Chilton, Didcot, 1974).
- [6] K. Stelle, Phys. Rev. **D16** (1977) 953; Gen. Rel. Grav. **9** (1978) 353; A. Salam and J. Strathdee, Phys. Rev. **D18** (1978) 4480.
- [7] J. Julve and M. Tonin, Nuovo Cim. **46B** (1978) 137.
- [8] E. Tomboulis, Phys. Lett. **70B** (1977) 361; Phys. Lett. **97B** (1980) 77; B. Hasslacher and E. Mottola, Phys. Lett. **99B** (1981) 221.
- [9] E. Fradkin and A. Tseytlin, Nucl. Phys. **B201** (1982) 469; Phys. Lett. **104B** (1981) 377.
- [10] E. Elizalde, S. Odintsov and A. Romeo, Phys. Rev. **D51** (1995) 1680, 4250.
- [11] M. Kaku, Nucl. Phys. **B203** (1982) 285; D. Boulware, G. Horowitz and A. Strominger, Phys. Rev. Lett. **50** (1983) 1726; E. Tomboulis, Phys. Rev. Lett. **52** (1984) 1173; Phys. Lett. **B389** (1996) 225.
- [12] R. Riegert, Phys. Lett. **134B** (1984) 56.
- [13] E. Fradkin and A. Tseytlin, Phys. Lett. **134B** (1984) 187.

- [14] I. Antoniadis and E. Mottola, Phys. Rev. **D45** (1992) 2013;
S. Odintsov, Z. Phys. **C54** (1992) 531.
- [15] I. Antoniadis, P. Mazur and E. Mottola, Nucl. Phys. **B388** (1992) 627.
- [16] I. Antoniadis, P. Mazur and E. Mottola, Phys. Lett. **B323** (1994) 284, **B394** (1997) 49;
I. Antoniadis and S. Odintsov, Phys. Lett. **B343** (1995) 76.
- [17] K. Hamada and F. Sugino, Nucl. Phys. **B553** (1999) 283, hep-th/9810095.
- [18] K. Hamada, Prog. Theor. Phys. **103** (2000) 1237, hep-th/9912098.
- [19] K. Hamada, Prog. Theor. Phys. **105** (2001) 673, hep-th/0012053.
- [20] T. Lee and G. Wick, Nucl. Phys. **B9** (1969) 209; Phys. Rev. **D2** (1970) 1033;
Phys. Rev. **D3** (1971) 1046;
R. Cutkosky, P. Landshoff, D. Olive and J. Polkinghorne, Nucl. Phys. **B12** (1969) 281;
N. Nakanishi, Prog. Theor. Phys. Suppl. **51** (1972) 1;
D. Boulware and D. Gross, Nucl. Phys. **B233** (1984) 1.
- [21] A. Polyakov, Phys. Lett. **103B** (1981) 207; Mod. Phys. Lett. **A2** (1987) 893.
- [22] V. Knizhnik, A. Polyakov and A. Zamolodchikov, Mod. Phys. Lett. **A3** (1988) 819.
- [23] J. Distler and H. Kawai, Nucl. Phys. **B321** (1989) 509;
F. David, Mod. Phys. Lett. **A3** (1988) 1651.
- [24] N. Seiberg, Prog. Theor. Phys. Suppl. **102** (1990) 319;
J. Polchinski, Proc. of String 1990 (Texas A& M, march 1990); Nucl. Phys. **B346** (1990) 253;
K. Hamada, Phys. Lett. **B324** (1994) 141; Nucl. Phys. **B413** (1994) 278.
- [25] K. Hamada, Phys. Lett. **B300** (1993) 322;
K. Hamada and A. Tsuchiya, Int. J. Mod. Phys. **A8** (1993) 4897.
- [26] P. Bouwknegt, J. McCarthy and K. Pilch, Commun. Math. Phys. **145** (1992) 541.
- [27] H. Kawai and M. Ninomiya, Nucl. Phys. **B336** (1990) 115;
H. Kawai, Y. Kitazawa and M. Ninomiya, Nucl. Phys. **B393** (1993) 280;
Nucl. Phys. **B404** (1993) 684;
T. Aida, Y. Kitazawa, H. Kawai and M. Ninomiya, Nucl. Phys. **B427** (1994) 158;
T. Aida, Y. Kitazawa, J. Nishimura and A. Tsuchiya, Nucl. Phys. **B444**

- (1995) 353;
H. Kawai, Y. Kitazawa and M. Ninomiya, Nucl. Phys. **B467** (1996) 313;
T. Aida and Y. Kitazawa, Nucl. Phys. **B491** (1997) 427.
- [28] S. Hathrell, Ann. Phys. **139** (1982) 136.
 - [29] S. Hathrell, Ann. Phys. **142** (1982) 34.
 - [30] M. Freeman, Ann. Phys. **153** (1984) 339.
 - [31] J. Wess and B. Zumino, Phys. Lett. **37B** (1971) 95.
 - [32] L. Bonora, P. Cotta-Ramusino and C. Reina, Phys. Lett. **B126** (1983) 305.
 - [33] D. Capper and M. Duff, Nuovo Cimento **23A** (1974) 173.
 - [34] S. Deser, M. Duff and C. Isham, Nucl. Phys. **B111** (1976) 45.
 - [35] M. Duff, Nucl. Phys. **B125** (1977) 334;
Twenty years of the Weyl anomaly, Class. Quant. Grav. **11** (1994) 1387,
hep-th/9308075, and references therein.
 - [36] K. Fujikawa, Phys. Rev. **D23** (1981) 2262; Phys. Rev. Lett. **44** (1980) 1733.
 - [37] S. Deser and A. Schwimmer, Phys. Lett. **B309** (1993) 279 hep-th/9302047.
 - [38] I. Drummond and S. Hathrell, Phys. Rev. **D21** (1980) 958.
 - [39] L. Abbott, Nucl. Phys. **B185** (1981) 189.
 - [40] E. DeRafael and J. Rosner, Ann. Phys. **82** (1974) 369.
 - [41] K. Hamada, Mod. Phys. Lett. **A16** (2001) 803, hep-th/0101100.
 - [42] D. Weingarten, Phys. Lett. **B90** (1980) 285, Nucl. Phys. **B210** [FS6] (1982) 229;
V. Kazakov, Phys. Lett. **150B** (1985) 282;
F. David, Nucl. Phys. **B257** (1985) 45;
J. Ambjørn, B. Durhuus and J. Fröhlich, Nucl. Phys. **B257** (1985) 433;
D. Boulatov, V. Kazakov, I. Kostov and A. Migdal, Nucl. Phys. **B275** (1986) 641.
 - [43] E. Brezin and V. Kazakov, Phys. Lett. **B236** (1990) 144;
M. Douglas and S. Shenker, Nucl. Phys. **B335** (1990) 635;
D. Gross and A. Migdal, Phys. Rev. Lett. **64** (1990) 127.
 - [44] J. Ambjørn, B. Durhuus and T. Jonsson, *Quantum Geometry*, (Cambridge Univ. Press, Cambridge, 1997).

- [45] M. Agishtein and A. Migdal, Nucl. Phys. **B385** (1992) 395;
 J. Ambjørn and J. Jurkiewicz, Phys. Lett. **B278** (1992) 42;
 S. Catterall, J. Kogut and R. Renken, Phys. Lett. **B328** (1994) 277.
- [46] S. Bilke, Z. Burda, A. Krzywicki, B. Petersson, J. Tabaczek and G. Thorleifsson, Phys. Lett. **B418** (1998) 266; Phys. Lett. **B432** (1998) 279.
- [47] H. Egawa, A. Fujitu, S. Horata, N. Tsuda and T. Yukawa, Nucl. Phys. **B (Proc. Suppl.) 73** (1999) 795, hep-lat/9809028;
 S. Horata, H. Egawa, N. Tsuda and T. Yukawa, Nucl. Phys. **B (Proc. Suppl.) 83** (2000) 751, hep-lat/9908048; Prog. Theor. Phys. **106** (2001) 1037, hep-lat/0004021;
 S. Horata, H. Egawa and T. Yukawa, Nucl. Phys. **B (Proc. Suppl.) 94** (2001) 701, hep-lat/0010050.
- [48] S. Horata, H. Egawa and T. Yukawa, Nucl. Phys. **B (Proc. Suppl.) 106** (2002) 971, hep-lat/0110042.
- [49] H. Hagura, Nucl. Phys. **B (Proc. Suppl.) 94** (2001) 704.